

Collins

# AQA GCSE Maths

Foundation Student Book  
Answers



William Collins' dream of knowledge for all began with the publication of his first book in 1819. A self-educated mill worker, he not only enriched millions of lives, but also founded a flourishing publishing house. Today, staying true to this spirit, Collins books are packed with inspiration, innovation and practical expertise. They place you at the centre of a world of possibility and give you exactly what you need to explore it.

Collins. Freedom to teach

---

Published by Collins  
An imprint of HarperCollins *Publishers*  
The News Building  
1 London Bridge Street  
London SE1 9GF

Browse the complete Collins catalogue at  
[www.collins.co.uk](http://www.collins.co.uk)

© HarperCollins *Publishers* Limited 2015

10 9 8 7 6 5 4 3 2 1

ISBN 978-0-00-814704-4

A catalogue record for this book is available from the British Library

Commissioned by Lucy Rowland and Katie Sergeant  
Project managed by Elektra Media and Hart McLeod Ltd  
Project edited by Caroline Green and Jennifer Yong  
Answers checked by Amanda Dickson and Steven Matchett  
Proofread by Helen Atkinson  
Illustrations by Ann Paganuzzi  
Cover design by We Are Laura  
Cover photographs by Godruma/Shutterstock (top) and Georgios Kollidas/Shutterstock (bottom)  
Production by Rachel Weaver

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the Publisher or a licence permitting restricted copying in the United Kingdom issued by the Copyright Licensing Agency Ltd., 90 Tottenham Court Road, London W1T 4LP.






#### **Acknowledgements**

The publishers gratefully acknowledge the permissions granted to reproduce copyright material in this book. Every effort has been made to contact the holders of copyright material, but if any have been inadvertently overlooked, the publisher will be pleased to make the necessary arrangements at the first opportunity.

## Chapter 1 – Number:

### Basic number

#### Exercise 1A

- 1 a 25, 62, 86, 151, 219, 310, 400, 501  
b 97, 357, 368, 740, 888, 2053, 4366  
c  $-9, -6, -2, 0, 2, 3, 4, 8$
- 2 a 700, 401, 174, 117, 80, 65, 18, 2  
b 6227, 3928, 2034, 762, 480, 395, 89, 59
- 3 a Number 4 (£128 250)  
b Number 1 (£129 100)  
c £850
- 4 a 368, 386, 638, 683, 836, 863  
b 368  
c 863
- 5 408, 480, 804, 840
- 6 33, 35, 38, 53, 55, 58, 83, 85, 88
- 7 7045 or 7405
- 8 a is smaller than    b is smaller than  
c is smaller than    d is bigger than  
e is smaller than    f is smaller than
- 9 a  $973 + 85 = 1058$  or  
 $975 + 83 = 1058$  or  
 $985 + 73 = 1058$  or  
 $983 + 75 = 1058$   
b  $357 - 98 = 259$
- 10 a is smaller than    b is bigger than  
c is smaller than    d is smaller than  
e is bigger than    f is smaller than  
g is smaller than    h is bigger than  
i is bigger than    j is smaller than  
k is smaller than    l is bigger than
- 11 a   
b   
c   
d 
- 12  $6^{\circ}\text{C}$      $-2^{\circ}\text{C}$      $-4^{\circ}\text{C}$      $2^{\circ}\text{C}$
- 13 
- 14 a 7, 8, 9, 10, 11  
b 6, 7, 8, 9, 10  
c 2, 3, 4, 5, 6, 7, 8

#### Exercise 1B

- 1 a 11    b 6    c 10    d 12  
e 11    f 13    g 11    h 12  
i 12    j 4    k 13    l 3
- 2 a 16    b 2    c 10    d 10  
e 6    f 18    g 6    h 15  
i 9    j 12    k 3    l 8
- 3 b  $3 + (2 \times 4) = 11$     c  $(9 \div 3) - 2 = 1$   
d  $9 - (4 \div 2) = 7$     e  $(5 \times 2) + 3 = 13$   
f  $5 + (2 \times 3) = 11$     g  $(10 \div 5) - 2 = 0$   
h  $10 - (4 \div 2) = 8$     i  $(4 \times 6) - 7 = 17$   
j  $7 + (4 \times 6) = 31$     k  $(6 \div 3) + 7 = 9$   
l  $7 + (6 \div 2) = 10$
- 4 a 38    b 48    c 3    d 2  
e 5    f 14    g 20    h 19  
i 54    j 7    k 2    l 7  
m 7    n 10    o 2
- 5 a  $(4 + 1)$   
b No brackets needed  
c  $(2 + 1)$   
d No brackets needed  
e  $(4 + 4)$   
f  $(16 - 4)$   
g No brackets needed  
h  $(5 + 5)$   
i  $(4 + 2)$   
j No brackets needed  
k No brackets needed  
l  $(8 - 2)$
- 6 a 8    b 6    c 13    d 9  
e 12    f 8    g 15  
h 16    i 1
- 7 No, correct answer is  $5 + 42 = 47$
- 8 a  $2 + 3 \times 4 = 14$   
b  $8 - 4 \div 4 = 7$  (correct)  
c  $6 + 3 \times 2 = 12$  (correct)  
d  $7 - 1 \times 5 = 2$   
e  $2 \times 7 + 2 = 16$  (correct)  
f  $9 - 3 \times 3 = 0$
- 9 a  $2 \times 3 + 5 = 11$     b  $2 \times (3 + 5) = 16$   
c  $2 + 3 \times 5 = 17$     d  $5 - (3 - 2) = 4$   
e  $5 \times 3 - 2 = 13$     f  $5 \times 3 \times 2 = 30$
- 10  $4 + 5 \times 3 = 19$   
 $(4 + 5) \times 3 = 27$ . So  $4 + 5 \times 3$  is smaller
- 11  $(5 - 2) \times 6 = 18$

12  $8 \div (5 - 3) = 4$

13  $10 - 3 \times 1.5$  and  $10 - 1.5 - 1.5 - 1.5$

- 14 a T            b F            c F  
 d T            e F            f T  
 g F            h F            i T  
 j T

15 a i  $(4 + 4) \div (4 + 4) = 1$

ii  $(4 \div 4) + (4 \div 4) = 2$

iii  $(4 + 4 + 4) \div 4 = 3$

iv  $(4 - 4) \times 4 + 4 = 4$

v  $(4 \times 4 + 4) \div 4 = 5$

b There are other ways to do these:

$4 + (4 + 4) \div 4 = 6$ ,  $4 + 4 - (4 \div 4) = 7$

$(4 + 4) \times (4 \div 4) = 8$ ,  $(4 + 4) + (4 \div 4) = 9$

c There are other ways to do this

$4 + \sqrt{4} + \sqrt{4} + \sqrt{4} = 10$

16

8	÷	4	+	6	=	8
×		+		-		
2	+	9	-	1	=	10
-		÷		×		
7	+	3	-	5	=	5
=	=	=				
9	7	1				

**Exercise 1C**

- 1 a 713            b 151            c 6381  
 d 968            e 622
- 2 a 646            b 826            c 3818  
 d 75.5            e 92.92            f 85.33
- 3 a 450            b 563            c 482  
 d 414            e 285            f 2892  
 g 4417            h 3767            i 4087  
 j 1828
- 4 a 128            b 29            c 334  
 d 17.8            e 61.63            f 6.882
- 5 a 558 miles    b 254 miles
- 6 252
- 7 Units digit should be 6 (from  $14 - 8$ )
- 8 For example:  $181 - 27 = 154$
- 9 a 115            b 204            c 294  
 d 212            e 425            f 150  
 g 800            h 960            i 1360  
 j 1518

- 10 a 294            b 370            c 288  
 d 83.2            e 21.63            f 25.2  
 g 16.44            h 321.5            i 30

- 11 a 219            b 317            c 315  
 d 10.6            e 0.99            f 12.1  
 g 25.2            h 1.41            i 1.44

12 a 705 miles    b £3525

- 13 a 47 miles  
 b Three numbers with a total of 125. First number must be less than 50, second number less than first number, third number less than second number, for example 48, 42 and 35.

- 14 a 119            b 96            c 144  
 d 210            e 210

- 15 a 13            b 37 weeks      c 4.3 m  
 d 36            e 45

- 16 a  $152 + 190 = 324$   
 b  $(10 \times 190) + 76 = 1976$   
 c  $(100 \times 38) + 190 = 3990$

**Exercise 1D**

- 1 a -1            b -1            c 2            d -3  
 e -4            f -6            g -6            h -1  
 i -5            j -4            k 4            l -1
- 2 a -4            b -4            c -10            d 2  
 e -32            f -30            g -8            h 10  
 i -36            j -14            k 12            l -101
- 3 a 6            b -5            c 6            d -1  
 e -2            f -6            g -9            h -5  
 i -80            j -7            k -1            l -47
- 4 a  $-5 + 7 = -2$   
 b  $+5 - 6 = -1$   
 c  $-3 - 8 = -11$
- 5 a 7 degrees            b  $-6^\circ\text{C}$
- 6 a  $2 - 8$   
 b  $2 + 5 - 8$  or  $2 + 4 - 7$  or  $8 - 4 - 5$  or  
 $8 - 2 - 7$  or  $5 - 4 - 2$   
 c  $2 - 5 - 7 - 8$             d  $2 + 5 - 4 - 7 - 8$
- 7 500 feet

**Exercise 1E**

- 1 a 6            b 7            c 8            d 6  
 e 8            f 10            g 2            h -3  
 i 2            j -3            k 1            l -5  
 m 3            n -4            o -3            p -8

2 a 7      b -8      c -5      d -11  
 e 11      f 6      g 8      h 8  
 i -2      j -1      k -9      l -5  
 m 5      n -9      o 8      p 0

3 a 3 °C      b 0 °C      c -3 °C  
 d -5 °C      e -11 °C

4 a 10 degrees Celsius  
 b 7 degrees Celsius  
 c 9 degrees Celsius

5 -9, -6, -5, -1, 1, 2, 3, 8

6 a 2      b -3      c -5  
 d -7      e -10      f -20

7 a 2      b 4      c -1  
 d -5      e -11      f 8

8 a 13      b 2      c 5  
 d 4      e 11      f -2

9 a -10      b -5      c -2  
 d 4      e 7      f -4

10 a -5      b 6      c 0  
 d -6      e -2      f 212  
 g -67      h 7      i 25

11 Yes. The worst case is at when the temperature is just below 0 °C and the thermometer error causes a measurement of  $0 + 2 = 2$  °C which is below 3 °C so the alert will come on.

12 a -1, 0, 1, 2, 3  
 b -9, -8, -7, -6, -5  
 c -2, -1, 0, 1, 2, 3; -4, -3, -2, -1, 0, 1  
 d -12, -11, -10, -9, -8, -7;  
 -14, -13, -12, -11, -10, -9  
 e -8, -7, -6, -5, -4, -3, -2;  
 -5, -4, -3, -2, -1, 0, 1  
 f -10, -9, -8, -7, -6, -5, -4; -1, 0, 1, 2, 3, 4, 5

13 a -4      b +3      c +4      d -6  
 e +7      f +2      g 0      h 0  
 i -6      j -7      k -9      l +4  
 m -3      n -9      o 0

14 a  $+6 + +5 = 11$       b  $+6 + -9 = -3$   
 c  $+6 - -9 = 15$       d  $+6 - +5 = 1$

15 a  $+5 + +7 - -9 = +21$   
 b  $+5 + -9 - +7 = -11$   
 c  $+7 + -7$  or  $+4 + -4$

16 -3 and 8

17  $-7\frac{1}{2}$  and  $-4\frac{1}{2}$

## Exercise 1F

1 a -15      b -14      c -24      d 6  
 e 14      f 2      g -2      h -8  
 i -4      j 3      k -24      l -10  
 m -18      n 16      o 36      p -4  
 q -12      r -4      s 7      t 25  
 u 18      v -8      w -45  
 x 3      y -40

2 a -9      b 16      c -3      d -32  
 e 18      f 18      g 6      h -4  
 i 20      j 16      k 8      l -48  
 m 13      n -13      o -8      p 0  
 q 16      r -42      s 6      t 1  
 u -14      v 6      w -4  
 x 7      y 0

3 a -2      b 30      c 15  
 d -27      e -7

4 a 4      b -9      c -3  
 d 6      e -4

5 a -9      b 3      c 1

6 a 16      b -2      c -12

7 a 24      b 6      c -4      d -2

8 For example:  $1 \times (-12)$ ,  $-1 \times 12$ ,  $2 \times (-6)$ ,  
 $6 \times (-2)$ ,  $3 \times (-4)$ ,  $4 \times (-3)$

9 For example:  $4 \div (-1)$ ,  $8 \div (-2)$ ,  $12 \div (-3)$ ,  
 $16 \div (-4)$ ,  $20 \div (-5)$ ,  $24 \div (-6)$

10 a 21      b -4      c 2      d -16  
 e 2      f -5      g -35      h -17  
 i -12      j 6      k 45      l -2  
 m 0      n -1      o -7      p -36  
 q 9      r 32      s 0      t -65

11 a -12      b 12 degrees      c  $3 \times -6$

12  $-5 \times 4$ ,  $3 \times -6$ ,  $-20 \div 2$ ,  $-16 \div -4$

13 a 16      b 41      c 21      d 64

## Exercise 1G

1 a 12 138      b 45 612      c 29 988  
 d 20 654      e 51 732      f 25 012  
 g 19 359      h 12 673      i 140 224  
 j 92 851      k 520 585      l 78 660

2 3500

3 No, 62 pupils cannot get in

4 Yes, he walks 57.6 km

- 5 Yes, 1 204 000 letters  
 6 5819 litres  
 7 No, she raised £302.40

### Exercise 1H

- 1 a 25                    b 15                    c 37  
 d 43                    e 27                    f 48  
 g 53                    h 52                    i 32  
 j 57                    k 37 rem 15        l 25 rem 5  
 m 34 rem 11        n 54 rem 9        o 36 rem 11  
 p 17 rem 4        q 23                    r 61 rem 14  
 s 42                    t 27 rem 2

- 2 68  
 3 38  
 4 4 months  
 5 34 hours  
 6 45p  
 7 a £46.75  
    b 13 weeks  
 8 An odd number divided by an even number cannot give a whole number  
 9 a 64                    b 22                    c 22

### Exercise 1I

- 1 a 89.28                b 298.39              c 66.04  
 d 167.98              e 2352.0              f 322.4  
 g 1117.8              h 4471.5              i 464.94  
 j 25.55                k 1047.2              l 1890.5  
 2 a £224.10        b £223.75        c £29.90  
 3  $5 \times 7 = 35$   
 4 £54.20  
 5 a £120.75        b  $£17 - £3.45 = £13.55$

### Review questions

- 1 a £1505  
    b 19 people  
 2 Murray did the addition first, Harry did the multiplication first.

- 3 a Adam has calculated  $5 \times 2$  instead of  $5^2$ , Bekki has added 3 to 5 first, instead of doing the power first.  
 b 26  
 4 a  $-2^\circ\text{C}$ ,  $-1^\circ\text{C}$ ,  $0^\circ\text{C}$ ,  $1^\circ\text{C}$   
 b 3 degrees  
 5 i Any number between  $-196^\circ\text{C}$  and  $-210^\circ\text{C}$ , e.g.  $-200^\circ\text{C}$   
    ii Any number over  $-196^\circ\text{C}$ , e.g.  $-10^\circ\text{C}$  or any positive number  
    iii Any number below  $-210^\circ\text{C}$ , e.g.  $-250^\circ\text{C}$   
 6 answer in the region of £7.50 to £8  
 7 DVD, jeans and pen  
 8 Pack of 8 at £0.625 each  
 9 a £49.90  
    b Small, so that all four children can have an ice cream (cost £4.80)  
 10 Yes. She only needed 8 paving stones.  
 11  $2.5 \times 8.4$ ,  $5 \times 4.2$ ,  $10 \times 2.1 = 21$   
 12 £133.05  
 13 a £141.75        b £119.25  
 14 a 7542  
    b 2457  
    c  $27 \times 2 = 54$

## Chapter 2 – Geometry and measures: Measures and scale drawings

### Exercise 2A

- 1 a 26 cm, 1.5 kg. Check individual answers.  
    b-j Check individual answers  
 2 The 5 metre ladder, since his height is about 175 cm and the lamp post will be about 525 cm.  
 3 Metres and centimetres are too small as units; needs to be a large unit as this is a large distance.  
 4 a 3400 mm    b 135 mm    c 67 cm  
    d 7030 m    e 2400 ml    f 590 cl  
    g 84 ml      h 5200 l      i 3750 kg  
    j 0.000 94 l    k 15 200 g    l 0.19 ml

- 5 a 1.25 m    b 8.2 cm    c 0.55 m  
 d 2.1 km    e 4.2 kg    f 5.75 t  
 g 2.58 l    h 3.4 l    i 0.6 t  
 j 63 cl    k 35 cm<sup>3</sup>    l 1.035 m<sup>3</sup>
- 6 He should choose the 2000 mm × 15 mm × 20 mm.
- 7 1 000 000
- 8 a 1 000 000  
 b 1 000 000 000 000

**Exercise 2B**

- 1 a 24 in    b 12 ft    c 3520 yd  
 d 80 oz    e 56 lb    f 6720 lb  
 g 40 pt    h 48 in    i 36 in  
 j 30 ft    k 64 oz    l 5 ft  
 m 70 lb    n 12 yd    o 224 oz
- 2 a 5 miles    b 120 pt    c 5280 ft  
 d 8 ft    e 7 st    f 7 gal  
 g 2 lb    h 5 yd    i 5 tons  
 j 63 360 in    k 8 lb    l 9 gal  
 m 10 st    n 3 miles    o 35 840 oz
- 3 The 32-ounce bag
- 4 4 014 489 600 in<sup>2</sup>
- 5 a miles, feet, ounces, gallons  
 b 108 inches  
 c 3 gallons

**Exercise 2C**

- 1 a 20 cm    b 13.2 lb    c 48 km  
 d 67.5 l    e 2850 ml    f 10 gal  
 g 12 in    h 50 miles    i 5 kg  
 j 3 pints    k 160 km    l 123.2 lb  
 m 180 l    n 90.9 kg    o 1100 yd  
 p 30 cm    q 6.4 kg    r 90 cm
- 2 ton
- 3 metre
- 4 a i 1000 g    ii 1 kg  
 b i 4500 g    ii 4.5 kg
- 5 a 135 miles    b 50 mph    c 2 h 42 min
- 6 4 hours 10 minutes
- 7 288 in<sup>3</sup>

**Exercise 2D**


- 1 a Onions: 40 m × 10 m; Soft fruits: 50 m × 10 m; Apple trees: 20 m × 20 m; Lawn: 30 m × 20 m; Potatoes: 50 m × 20 m  
 b Onions: 400 m<sup>2</sup>; Soft fruit: 500 m<sup>2</sup>; Apple trees: 400 m<sup>2</sup>; Lawn: 600 m<sup>2</sup>; Potatoes: 1000 m<sup>2</sup>
- 2 a 33 cm    b 9 cm
- 3 a 6 m × 6 m    b 8 m × 2 m    c 4 m × 3 m  
 d 4 m × 6 m    e 6 m × 4 m    f 3 m × 2 m
- 4 a Student's scale drawing  
 b 39 plants (the edge is about 19 m)
- 5 a 15 km    b 9 km    c 12 km  
 d 13.5 km    e 7.8 km    f 5.4 km
- 6 a On the map, Carlisle to Scarborough is 5.8cm, so 1cm represents  $232 \times 100000 / 5.8 = 4000000$   
 b i 176 km    ii 66 km
- 7 a 80 km    b 70 km    c 122 km
- 8 d 1:600
- 9 a i 160 miles    ii 310 miles

**Exercise 2E**




- 1 Bicycle about 2 m, bus about 10 m, train about 17 m
- 2 Height about 4 m, length about 18 or 19 m
- 3 a About 4 m  
 b About 5 m  
 c About 5.5 m
- 4 About 9 m; the ratio of Joel's height in the photograph to his real height must be the same as the ratio of the height on the statue in the photograph to its actual height
- 5 Assuming the man's height is about 1.8 metres, the T. Rex height is about 4 times the height of the man and  $4 \times 1.8 = 7.2\text{m}$ . So yes, Ben's estimate of the height is correct. The length of T. Rex looks about 5 times the height of the man.  $5 \times 1.8 = 9\text{m}$ , so Ben has overestimated the length of the T. Rex, but not by much.

**Exercise 2F**

1 A, C, D






3 Yes. For example, Jenny could make it from a card  $10\text{ cm} \times 15\text{ cm}$




7 a and b

8 The length and height of the net needs to be at least  $5.45\text{ cm} + 5\text{ cm} + 5.45\text{ cm} = 15.9\text{ cm}$ , so a card  $15\text{ cm}$  by  $15\text{ cm}$  is too small.

**Exercise 2G**





4 Students' own diagram





**b** She could be correct as 4 bricks could be put together to form a 2 cm by 12 cm block, 4 such blocks creates the bottom layer of bricks almost filling the base of  $12\text{ cm} \times 9\text{ cm}$ , making  $4 \times 4 = 16$  bricks in a layer. There will be 5 such layers, making  $5 \times 16 = 80$  bricks, leaving a  $1\text{ cm} \times 12\text{ cm} \times 5\text{ cm}$  gap at the end of the box. You can put two of the blocks sideways into this space, giving 8 extra bricks and a total of 88 bricks.



## Review questions

1

The length of a garden	metres
The mass of a puppy	kilograms
The length of a brick	centimetres

2 **a** 40 000 metres **b** 25 miles

3 **a** **i** Kilometres **ii** Litres


**b** **i** 90 mm **ii** 3 kg

4 Andrew is correct as  $130 \times \frac{5}{8} = 81.25$

5 **a** 1 m **b** 2.82 m

6 **a** 160 mm **b** 150 m

7




8 Centimetres is too small and this distance is an approximation of a large distance and so needs to be a large unit such as kilometres.


9 **a** Seven and a half miles

**b** 2.3 kg potatoes, 23 grams of butter, 2.3 litres of milk

10



11



## Chapter 3 – Statistics: Charts, tables and averages

### Exercise 3A

1 a

Goals	0	1	2	3
Frequency	6	8	4	2

b 1 goal      c 22

2 a

Score	1	2	3	4	5	6
Frequency	5	6	6	6	3	4

b 30      c Yes, frequencies are similar.

3 a/b

Temperature (°C)	Frequency
14–16	5
17–19	10
20–22	8
23–25	5
26–28	2

c 17–19 °C

d Getting warmer in the first half of the month and then getting cooler towards the end.

4 a/b

Height(cm)	Frequency
151–155	2
156–160	5
161–165	5
166–170	7
171–175	5
176–180	4
181–185	3
186–190	1

c 166–170 cm


- 5 a Observation      b Sampling  
 c Observation      d Sampling  
 e Experiment      f Observation  
 g Experiment

6 Draw a tally chart with class intervals 1–10, 11–20, 21–30, 31–40, 41–50 and 51–60. Then complete a grouped frequency table from the tally chart.

7 The class intervals overlap as the ages 20, 30, 40 and 50 are in two different class intervals

### Exercise 3B

1



2 a May 9 h, Jun 11 h, Jul 12 h, Aug 11 h, Sep 10 h


b July      c Visual impact, easy to understand.

3 a Simon      b £330  
 c Difficult to show fractions of a £ symbol.

4 a Swimming      b 74  
 c a better visual impact and easier to read the frequencies


d No. It may not include people who are not fit or not interested in sport.


5 a



Key = 1 accident

b






**d** Use the pictogram because an appropriate symbol makes more impact.

**6 a**


Time (min)	Frequency
1-10	4
11-20	7
21-30	5
31-40	5
41-50	
51-60	2



**c** Some students live close to the school and some live a good distance away and probably travel to school by bus.

**7** Yes, in most cities. If you double the minimum temperature each time, it is very close to the maximum temperature.

**8** The bar chart is misleading because the temperature axis does not start at zero.




**10 a** 102      **b** 16      **c** 42

**11** £88.80

**Exercise 3C**

**1 a** Tuesday, 52p      **b** 2p  
**c** Friday      **d** £90




**b** about 16 500

**c** 1991 and 2001


**d** No; do not know the reason why the population started to decrease after 2001

**3** All the temperatures were higher than 20 degrees.




**b** 112

5 a




- b About 60 million  
 c 1985 and 1990  
 d e.g. the trend in tourism is rising over the 30 year period. This is probably because holidays have become cheaper and the rise in popularity of package holidays

6 a



b 7 °C and 10 °C

7 a




b It is not possible to predict the trend as the puppy's mass is likely to level out.

**Exercise 3D**

- 1 a 4      b 48      c -1  
 d  $\frac{1}{4}$       e no mode      f 3.21  
 2 a red      b sun      c  $\beta$       d M  
 3 a 32      b 6      c no  
 d no; boys generally take larger shoe sizes  
 4 a 5  
 b no; more than half the class got a higher mark

5 a



- b 70  
 c 24  
 d cannot tell; know only that 9 households had between 0 and 4 emails  
 e 10-14

6 a 28

- b i brown      ii blue      iii brown  
 c Both students had blue eyes.

7 a May lose count.

- b Put in a table, or arrange in order  
 c 4

8 The mode will be the most popular item or brand sold in a shop.

**Exercise 3E**

- 1 a 5      b 33      c  $7\frac{1}{2}$       d 24  
 e  $8\frac{1}{2}$       f 0      g 5.25  
 2 a £2.20      b £2.25  
 c median, because it is the central value

- 3 a 5  
b i 5      ii 215      iii 10      iv 10

- 4 a 13 years, Bella  
b 162 cm, Dolly  
c 46 kg, Amina  
d Bella, because she is closest to the 3 medians

- 5 A median of £15 does not take into account the large payment of the £200 so is in no way representative.

- 6 a Possible answer: 11, 15, 21, 21 (one below or equal to 12 and three above or equal)

b Any four numbers higher than or equal to 12, and any two lower or equal to 12

c Eight, all 4 or under

- 7 12, 14, 14, 16, 20, 22, 24, or 12, 14, 14, 16, 18, 20, 24 or 12, 14, 14, 16, 18, 22, 24

- 8 a 21      b 16

c

Mark	Frequency
12	1
13	3
14	4
15	3
16	6
17	3
18	1

- d There are 21 students altogether, so the median is the 11<sup>th</sup> mark when the marks are put in order.

There are 11 values up to a mark of 15, so the median is 15.

### Exercise 3F

- 1 a 6      b 24      c 45  
d 1.57      e 2

- 2 a 55.1      b 324.7      c 58.5  
d 44.9      e 2.3

- 3 a 61      b 60      c 59  
d Blake      e 2

- 4 42 min

- 5 a £200      b £260      c £278  
d Median, because the extreme value of £480 is taken into account when calculating the mean and the mode would be too small to represent the data.

- 6 a 35      b 36

- 7 a 6  
b 16; all the numbers and the mean are 10 more than those in part a  
c i 56      ii 106      iii 7

- 8 Possible answers: Speed – Kath, James, John, Joseph; Roberts – Frank, Jim, Helen, Evie.

- 9 36

- 10 24

### Exercise 3G

- 1 a i Mode 3, median 4, mean 5

ii 6, 7,  $7\frac{1}{2}$       iii 4, 6, 8

b i Mean: balanced data

ii Mode: 6 appears five times

iii Median: 28 is an extreme value

- 2 a i 29      ii 28      iii 27.1

b The mean as it is balanced data

- 3 a Mode 73, median 76, mean 80

b The mean, because it is the highest average

- 4 a Mean – takes all marks into account

b Median – avoids any extreme amounts

c Mode – non numerical data

d Median – avoids any extreme heights

e Mode – non numerical data

f Mean – takes all masses into account

- 5 Yes, mode is 31, median is 31, and mean is  $31\frac{1}{2}$ ; all are close to the stated average.

- 6 a i £20 000      ii £28 000      iii £34 000

b i The 6% rise, because it gives a greater increase in salary for the higher paid employees.

ii 6% increase: £21 200, £29 680, £36 040; £1500 pay rise: £21 500, £29 500, £35 500

- 7 a Median      b Mode      c Mean

- 8 Tom – mean, takes all scores into account; David – median, avoids the extreme scores; Mohammed – mode, the two highest scores


- 9 The boss chose the mean while the worker chose the mode.

### Exercise 3H

- 1 a 7                      b 26                      c 5  
d 2.4                      e 7
- 2 a  $5^\circ, 3^\circ, 2^\circ, 7^\circ, 3^\circ$   
b Variable or inconsistent weather over England
- 3 a £31, £28, £33                      b £8, £14, £4  
c Not particularly consistent
- 4 a 82 and 83                      b 20 and 12  
c Shona, because the ranges show that her scores are more consistent
- 5 a i 5 min and 4 min  
ii 9 min and 13 min  
b Number 50, because the times are more consistent
- 6 a Isaac, Oliver, Andrew, Niamh, Zoe, Jake and Zac  
b For example: 70 cm to 90 cm
- 7 a The teachers because they have a high mean and students could not have a range of 20.  
b Year 11 students as the mean is between 15 and 16 and the range is 1.
- 8 For example: a 1, 6, 6, 6, 6    b 2, 5, 5, 6, 7

c on average Josh spent more time on his computer, but Hannah's times were more consistent.

- 3 a 84 cm    b 73 cm    c 70 cm  
d the median, avoids the extreme height of 43 cm
- 4 The modal age of 16, as most of the team are this age




- 6 The vertical scale does not start at zero, the horizontal scale is not linear, no £ sign for sales, no information to explain what it is about
- 7 a 12                      b 11                      c 11
- 8 11.6

### Review questions

1 a

Number		Frequency
1		3
2		4
3		7
4		5
5		4
6		1



- 2 a i 75 minutes                      ii 80 minutes  
b i 110 minutes                      ii 140 minutes

## Chapter 4 – Geometry and measures: Angles

### Exercise 4A

- 1 a  $48^\circ$                       b  $307^\circ$                       c  $108^\circ$   
d  $52^\circ$                       e  $59^\circ$                       f  $81^\circ$
- 2 a  $82^\circ$                       b  $105^\circ$                       c  $75^\circ$
- 3 No, as  $45^\circ + 125^\circ = 170^\circ$  and for a straight line it should be  $180^\circ$
- 4 a  $x = 100^\circ$     b  $x = 110^\circ$     c  $x = 30^\circ$
- 5 a  $x = 55^\circ$     b  $x = 45^\circ$     c  $x = 12.5^\circ$
- 6 a  $x = 34^\circ, y = 98^\circ$                       b  $x = 70^\circ, y = 120^\circ$   
c  $x = 20^\circ, y = 80^\circ$
- 7  $6 \times 60^\circ = 360^\circ$ ; imagine six of the triangles meeting at a point
- 8  $x = 35^\circ, y = 75^\circ$ ;  $2x = 70^\circ$  (opposite angles), so  $x = 35^\circ$  and  $x + y = 110^\circ$  (angles on a line), so  $y = 75^\circ$

## Exercise 4B

- 1 a  $70^\circ$     b  $50^\circ$     c  $80^\circ$     d  $60^\circ$   
e  $75^\circ$     f  $109^\circ$     g  $38^\circ$     h  $63^\circ$
- 2 a No, total is  $190^\circ$     b Yes, total is  $180^\circ$   
c No, total is  $170^\circ$     d Yes, total is  $180^\circ$   
e Yes, total is  $180^\circ$     f No, total is  $170^\circ$
- 3 a  $80^\circ$     b  $67^\circ$     c  $20^\circ$   
d  $43^\circ$     e  $10^\circ$     f  $1^\circ$
- 4 a–c Students' own drawings    d  $180^\circ$
- 5 a  $60^\circ$     b Equilateral triangle  
c Same length
- 6 a  $70^\circ$  each    b Isosceles triangle  
c Same length
- 7 a  $109^\circ$     b  $130^\circ$     c  $135^\circ$
- 8  $65^\circ$
- 9 Joe is not correct as  $\text{DFE} = 30^\circ$ ,  $\text{DEF} = 75^\circ$   
hence angle D =  $180^\circ - 105^\circ = 75^\circ$  but  
Hannah is correct as  $\text{FED} = \text{FDE} = 75^\circ$
- 10  $a = 35^\circ$  (angles in a triangle) because the  
other angles in the triangle are  $65^\circ$  (angles  
on a line) and  $80^\circ$  (opposite angles) giving  
a total of 145, this subtracted from the 180  
degrees in a triangle leaves the answer of  
35. As  $a$  is less than  $90^\circ$ , it is acute.
- 11 Missing angle =  $y$ ,  $x + y = 180^\circ$  and  
 $a + b + y = 180^\circ$  so  $x = a + b$
- 12  $32^\circ$
- 13  $72^\circ$

## Exercise 4C

- 1c  $2, 2 \times 180^\circ = 360^\circ$
- 2c  $3, 3 \times 180^\circ = 540^\circ$
- 3c  $4, 4 \times 180^\circ = 720^\circ$

4

Shape	Number of sides	Triangles	Angle sum
triangle	3	1	$180^\circ$
quadrilateral	4	2	$360^\circ$
pentagon	5	3	$540^\circ$
hexagon	6	4	$720^\circ$
heptagon	7	5	$900^\circ$
octagon	8	6	$1080^\circ$
nonagon	9	7	$1260^\circ$
decagon	10	8	$1440^\circ$

- 5  $18, 18 \times 180^\circ = 3240^\circ$

## Exercise 4D

- 1 a  $90^\circ$     b  $150^\circ$     c  $80^\circ$   
d  $80^\circ$     e  $77^\circ$     f  $131^\circ$
- 2 a No, total is  $350^\circ$     b Yes, total is  $360^\circ$   
c No, total is  $350^\circ$     d No, total is  $370^\circ$   
e Yes, total is  $360^\circ$     f Yes, total is  $360^\circ$
- 3 a  $100^\circ$     b  $67^\circ$     c  $120^\circ$   
d  $40^\circ$     e  $40^\circ$     f  $1^\circ$
- 4 a  $90^\circ$     b Rectangle    c Yes, square
- 5 a  $120^\circ$     b  $170^\circ$     c  $125^\circ$   
d  $136^\circ$     e  $149^\circ$     f  $126^\circ$
- 6  $60^\circ + 60^\circ + 120^\circ + 120^\circ + 120^\circ + 240^\circ = 720^\circ$
- 7  $y = 360^\circ - 4x$ ;  $2x + y + 2x = 360^\circ$ ,  $4x + y = 360^\circ$ ,  
so  $y = 360^\circ - 4x$
- 8  $x = 40^\circ$ , smallest angle =  $60^\circ$

## Exercise 4E

1

Regular polygon	Number of sides	Interior angle sum	Each interior angle
octagon	8	$1080^\circ$	$135^\circ$
nonagon	9	$1260^\circ$	$140^\circ$
decagon	10	$1440^\circ$	$144^\circ$

2

Regular polygon	Number of sides	Interior angle	Exterior angle
square	4	$90^\circ$	$90^\circ$
pentagon	5	$108^\circ$	$72^\circ$
hexagon	6	$120^\circ$	$60^\circ$
octagon	8	$135^\circ$	$45^\circ$
nonagon	9	$140^\circ$	$40^\circ$
decagon	10	$144^\circ$	$36^\circ$

- 3 a i  $45^\circ$     ii 8    iii  $1080^\circ$   
b i  $20^\circ$     ii 18    iii  $2880^\circ$   
c i  $15^\circ$     ii 24    iii  $3960^\circ$   
d i  $36^\circ$     ii 10    iii  $1440^\circ$
- 4 a i  $172^\circ$     ii 45    iii  $7740^\circ$   
b i  $174^\circ$     ii 60    iii  $10\,440^\circ$   
c i  $156^\circ$     ii 15    iii  $2340^\circ$   
d i  $177^\circ$     ii 120    iii  $21\,240^\circ$
- 5 a Exterior angle is  $7^\circ$ , which does not  
divide exactly into  $360^\circ$   
b Exterior angle is  $19^\circ$ , which does not  
divide exactly into  $360^\circ$

- c Exterior angle is  $11^\circ$ , which does divide exactly into  $360^\circ$   
 d Exterior angle is  $70^\circ$ , which does not divide exactly into  $360^\circ$
- 6 a  $7^\circ$  does not divide exactly into  $360^\circ$   
 b  $26^\circ$  does not divide exactly into  $360^\circ$   
 c  $44^\circ$  does not divide exactly into  $360^\circ$   
 d  $13^\circ$  does not divide exactly into  $360^\circ$
- 7  $x = 45^\circ$ , they are the same, true for all regular polygons
- 8 Three are  $135^\circ$  and two are  $67.5^\circ$
- 9  $48^\circ$ ;  $\frac{1440^\circ - 5 \times 240^\circ}{5}$
- 10 10
- 11  $72^\circ, 72^\circ, 108^\circ, 144^\circ, 144^\circ$
- 12  $93^\circ$

### Exercise 4F


- 1 a  $d$                       b  $f$                       c  $d$   
 d  $f$                       e  $f$                       f  $e$
- 2 a  $40^\circ$   
 b  $b = c = 70^\circ$   
 c  $d = 75^\circ, e = f = 105^\circ$   
 d  $g = 50^\circ, h = i = 130^\circ$   
 e  $j = k = l = 70^\circ$   
 f  $n = m = 80^\circ$   
 g  $a = 50^\circ, b = 130^\circ$   
 h  $c = d = 65^\circ, e = f = 115^\circ$   
 i  $g = i = 65^\circ, h = 115^\circ$   
 j  $j = k = 72^\circ, l = 108^\circ$   
 k  $m = n = o = p = 105^\circ$   
 l  $q = r = s = 125^\circ$
- 3 a  $a = 95^\circ$       b  $b = 66^\circ, c = 114^\circ$
- 4 a  $x = 30^\circ, y = 120^\circ$       b  $x = 25^\circ, y = 105^\circ$   
 c  $x = 30^\circ, y = 100^\circ$       d  $x = 50^\circ, y = 110^\circ$   
 e  $x = 25^\circ, y = 55^\circ$       f  $x = 20^\circ, y = 140^\circ$
- 5  $290^\circ$ ;  $x$  is double the angle allied to  $35^\circ$ , so is  $2 \times 145^\circ$
- 6  $a = 66^\circ$ ; angle  $BDC = 66^\circ$  (angles in a triangle =  $180^\circ$ ) and  $a$  is the corresponding angle
- 7 Angle  $PQD = 64^\circ$  (alternate angles), so angle  $DQY = 116^\circ$  (angles on a line =  $180^\circ$ )
- 8 Use alternate angles to see  $b, a$  and  $c$  are all angles on a straight line, and so total  $180^\circ$
- 9 Third angle in triangle equals  $q$  (alternate angle), angle sum of triangle is  $180^\circ$

### Exercise 4G

- 1 a  $a = 110^\circ, b = 55^\circ$       b  $c = 75^\circ, d = 115^\circ$   
 c  $e = 87^\circ, f = 48^\circ$
- 2 a  $a = c = 105^\circ, b = 75^\circ$       b  $d = f = 70^\circ, e = 110^\circ$   
 c  $g = i = 63^\circ, h = 117^\circ$
- 3 a  $a = c = 105^\circ, b = 75^\circ$       b  $d = f = 93^\circ, e = 87^\circ$   
 c  $g = i = 49^\circ, h = 131^\circ$
- 4 a  $a = 135^\circ, b = 25^\circ$       b  $c = d = 145^\circ$   
 c  $e = f = 94^\circ$
- 5 a  $a = 58^\circ, b = 47^\circ$       b  $c = 141^\circ, d = 37^\circ$   
 c  $e = g = 65^\circ, f = 115^\circ$
- 6 a  $65^\circ$       b  $60^\circ$       c  $68^\circ$
- 7 both  $129^\circ$
- 8 Marie is correct as a rectangle is a special form of parallelogram with all angles =  $90^\circ$
- 9 a  $65^\circ$   
 b Trapezium, angle  $A +$  angle  $D = 180^\circ$  and angle  $B +$  angle  $C = 180^\circ$

10  $135^\circ$

11



- 12 A trapezium; angles add up to  $10x$ , two angles  $x$  and  $4x = 2x + 3x$ , the other pair of angles. Hence each pair adds up to 180 (since  $2 \times 180 = 360$ ). Hence two pairs of allied angles, hence a trapezium. Alternatively you could have found that  $x = 36$  which will give the same result.

### Exercise 4H

- 1 a  $110^\circ, 12.6$  km      b  $250^\circ, 4.5$  km  
 c  $091^\circ, 11.8$  km      d  $270^\circ, 8.4$  km  
 e  $130^\circ, 7.2$  km      f  $180^\circ, 4.2$  km
- 2 Students' sketches
- 3 a Sketch  
 b D is due south of B and B is east of A, so A must be west of D. A bearing to the west will be greater than  $180^\circ$
- 4 a  $090^\circ, 180^\circ, 270^\circ$       b  $000^\circ, 270^\circ, 180^\circ$
- 5 a  $250^\circ$       b  $325^\circ$       c  $144^\circ$



- 6 a  $045^\circ$       b  $286^\circ$   
 c Measure the distance from X to Y and divide 15 by this to find the scale of the map. Then measure the distance from Y to P and multiply by the scale factor.
- 7 a 900 m      b  $280^\circ$   
 c angle NHS =  $150^\circ$  and HS = 3 cm
- 8  $108^\circ$
- 9  $255^\circ$
- 10 9.2 km

## Review questions

- 1  $16^\circ$
- 2 a i  $115^\circ$   
 ii opposite angles are equal and the angle opposite to  $x$  is  $180 - 65$   
 b the angles do not add up to  $360^\circ$
- 3 a  $50^\circ$       b  $32.5^\circ$
- 4  $3 \times 180 = 540^\circ$  A five-sided shape can be split into three triangles hence  $3 \times 180$
- 5  $150^\circ$
- 6 angle TQP =  $37^\circ$ , alternate angles, PTQ =  $180 - (29 + 37) = 114$  angles in a triangle, QTS =  $180 - 114$ , angles on a line, = 66.
- 7  $333^\circ$
- 8  $360 \div 8 = 45^\circ$ , exterior angle formula is  $\frac{360}{\text{number of slides}}$ , in this case
- 9  $180 - (360 \div 6) = 120^\circ$
- 10 Selvi might be correct. You will need to draw one example showing this is not a kite, and one example showing that this could be a kite
- 11 b 12.4 km

## Chapter 5 – Number: Number properties

### Exercise 5A

- 1 a 3, 6, 9, 12, 15      b 7, 14, 21, 28, 35  
 c 9, 18, 27, 36, 45      d 11, 22, 33, 44, 55  
 e 16, 32, 48, 64, 80
- 2 a 254, 108, 68, 162, 98, 812, 102, 270  
 b 111, 255, 108, 162, 711, 615, 102, 75, 270  
 c 255, 615, 75, 270      d 108, 162, 711, 270

- 3 a 72, 132, 216, 312, 168, 144  
 b 161, 91, 168, 294  
 c 72, 102, 132, 78, 216, 312, 168, 144, 294
- 4 a 98      b 99      c 96  
 d 95      e 98      f 96
- 5 a 1002      b 1008      c 1008
- 6 No, 50 is not a multiple of 6.
- 7 4 or 5 (as 2, 10 and 20 are not realistic answers)
- 8 a 18      b 28      c 15
- 9 66
- 10 5 numbers: 18, 36, 54, 72, 90
- 11 a 1, 2, 5, 10  
 b 1, 2, 3, 5, 6, 10, 15, 50  
 c 1, 2, 4, 5, 10, 20, 25, 50, 100  
 d For example, 60, 72, 84, 96

### Exercise 5B

- 1 a 1, 2, 5, 10      b 1, 2, 4, 7, 14, 28  
 c 1, 2, 3, 6, 9, 18      d 1, 17  
 e 1, 5, 25      f 1, 2, 4, 5, 8, 10, 20, 40  
 g 1, 2, 3, 5, 6, 10, 15, 30  
 h 1, 3, 5, 9, 15, 45      i 1, 2, 3, 4, 6, 8, 12, 24  
 j 1, 2, 4, 8, 16
- 2 8 ways (1, 2, 3, 4, 6, 8, 12, 24 per box)
- 3 a 1, 2, 3, 4, 5, 6, 8, 10, 12, 15, 20, 24, 30, 40, 60, 120  
 b 1, 2, 3, 5, 6, 10, 15, 25, 30, 50, 75, 150  
 c 1, 2, 3, 4, 6, 8, 9, 12, 16, 18, 24, 36, 48, 72, 144  
 d 1, 2, 3, 4, 5, 6, 9, 10, 12, 15, 18, 20, 30, 36, 45, 60, 90, 180  
 e 1, 13, 169  
 f 1, 2, 3, 4, 6, 9, 12, 18, 27, 36, 54, 108  
 g 1, 2, 4, 7, 14, 28, 49, 98, 196  
 h 1, 3, 9, 17, 51, 153  
 i 1, 2, 3, 6, 9, 11, 18, 22, 33, 66, 99, 198  
 j 1, 199
- 4 a 55      b 67      c 29      d 39  
 e 65      f 80      g 80      h 70  
 i 81      j 50
- 5 a 2      b 2      c 3      d 5  
 e 3      f 3      g 7      h 5  
 i 10      j 11
- 6 4 in each layer, which is a  $2 \times 2$  square
- 7 It does not have a factor of 3.
- 8 5

### Exercise 5C

- 23 and 29
- 97
- All these numbers are not prime.
- 3, 5, 7
- only if all 31 bars are in a single row, as 31 is a prime number and its only factors are 1 and 31
- a E                    b C                    c C
- a as this would leave 27 and taking off any other prime, which must be odd, would leave an even number  
b 3, 7 and 19 or 5, 7 and 17 or 5, 11 and 13
- All prime numbers greater than 2 are odd and Odd – Odd = Even.
- $97 - 61 = 36$ ,  $89 - 73 = 16$ ,  $89 - 53 = 36$ ,  $83 - 67 = 16$ ,  $83 - 47 = 36$ ,  $83 - 19 = 64$ ,  $79 - 43 = 36$ ,  $73 - 37 = 36$ ,  $67 - 31 = 36$ ,  $59 - 43 = 16$ ,  $59 - 23 = 36$ ,  $53 - 37 = 16$ ,  $53 - 17 = 36$ ,  $47 - 31 = 16$ ,  $47 - 11 = 36$ ,  $29 - 13 = 16$

### Exercise 5D

- a  $84 = 2 \times 2 \times 3 \times 7$   
b  $100 = 2 \times 2 \times 5 \times 5$   
c  $180 = 2 \times 2 \times 3 \times 3 \times 5$   
d  $220 = 2 \times 2 \times 5 \times 11$   
e  $280 = 2 \times 2 \times 2 \times 5 \times 7$   
f  $128 = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2$   
g  $50 = 2 \times 5 \times 5$   
h  $1000 = 2 \times 2 \times 2 \times 5 \times 5 \times 5$   
i  $576 = 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 3 \times 3$   
j  $650 = 2 \times 5 \times 5 \times 13$
- a  $2^2 \times 3 \times 7$                     b  $2^2 \times 5^2$   
c  $2^2 \times 3^2 \times 5$                     d  $2^2 \times 5 \times 11$   
e  $2^3 \times 5 \times 7$                     f  $2^7$   
g  $2 \times 5^2$                     h  $2^3 \times 5^3$   
i  $2^6 \times 3^2$                     j  $2 \times 5^2 \times 13$
- 1, 2, 3,  $2^2$ , 5,  $2 \times 3$ , 7,  $2^3$ ,  $3^2$ ,  $2 \times 5$ , 11,  $2^2 \times 3$ , 13,  $2 \times 7$ ,  $3 \times 5$ ,  $2^4$ , 17,  $2 \times 3^2$ , 19,  $2^2 \times 5$ ,  $3 \times 7$ ,  $2 \times 11$ , 23,  $2^3 \times 3$ ,  $5^2$ ,  $2 \times 13$ ,  $3^3$ ,  $2^2 \times 7$ , 29,  $2 \times 3 \times 5$ , 31,  $2^5$ ,  $3 \times 11$ ,  $2 \times 17$ ,  $5 \times 7$ ,  $2^2 \times 3^2$ , 37,  $2 \times 19$ ,  $3 \times 13$ ,  $2^3 \times 5$ , 41,  $2 \times 3 \times 7$ , 43,  $2^2 \times 11$ ,  $3^2 \times 5$ ,  $2 \times 23$ , 47,  $2^4 \times 3$ ,  $7^2$ ,  $2 \times 5^2$
- a Each is double the previous number  
b 64, 128  
c 81, 243, 729  
d 256, 1024, 4096  
e  $3, 3^2, 3^3, 3^4, 3^5, 3^6, \dots; 4, 4^2, 4^3, 4^4, 4^5, \dots$

- a  $2 \times 2 \times 3 \times 5$   
b  $2^2 \times 3 \times 5$   
c  $120 = 2^3 \times 3 \times 5$ ,  $240 = 2^4 \times 3 \times 5$ ,  $480 = 2^5 \times 3 \times 5$
- a  $7^2 \times 11^2 \times 13^2$   
b  $7^3 \times 11^3 \times 13^3$   
c  $7^{10} \times 11^{10} \times 13^{10}$
- because 3 is not a factor of 40 so it does not divide exactly
- a  $5 \times 7 \times 11$                     b  $3 \times 7^2$ ,  $x = 3$ ,  $y = 7$

### Exercise 5E

- a 20                    b 56                    c 6                    d 28  
e 10                    f 15                    g 24                    h 30
- It is their product.
- a 8                    b 18                    c 12                    d 30
- No. Because the numbers in each part have common factors
- a 168                    b 105                    c 84                    d 84  
e 96                    f 54                    g 75                    h 144
- 3 packs of cheese slices and 4 packs of bread rolls
- a 8                    b 7                    c 4                    d 14  
e 4                    f 9                    g 5                    h 4  
i 3                    j 16                    k 5                    l 18
- a i no                    ii yes                    iii yes                    iv no  
b i no                    ii no                    iii yes                    iv no
- 18 and 24, or 6 and 72

### Exercise 5F

- 36, 49, 64, 81, 100, 121, 144, 169, 196, 225, 256, 289, 324, 361, 400
- 4, 9, 16  
 $1 + 3 + 5 + 7 + 9 = 25$   
 $1 + 3 + 5 + 7 + 9 + 11 = 36$   
 $1 + 3 + 5 + 7 + 9 + 11 + 13 = 49$
- a 3                    b 5                    c 7  
d odd numbers
- a 529                    b 3249                    c 5929  
d 15 129                    e 23 104                    f 10.24  
g 90.25                    h 566.44                    i 16  
j 144
- a 169                    b 196  
c Answer between 169 and 196 (exact answer is 174.24)

- 6 a 50, 65, 82      b 98, 128, 162  
 c 51, 66, 83      d 48, 63, 80  
 e 149, 164, 181
- 7 a 25, 169, 625, 1681, 3721  
 b Answers in each row are the same
- 8 £144
- 9 29
- 10 36 and 49

**Exercise 5G**

- 1 a 6, 12, 18, 24, 30      b 13, 26, 39, 52, 65  
 c 8, 16, 24, 32, 40      d 20, 40, 60, 80, 100  
 e 18, 36, 54, 72, 90
- 2 1, 4, 9, 16, 25, 36, 49, 64, 81, 100
- 3 a 1, 2, 3, 4, 6, 12      b 1, 2, 4, 5, 10, 20  
 c 1, 3, 9      d 1, 2, 4, 8, 16, 32  
 e 1, 2, 3, 4, 6, 8, 12, 24      f 1, 2, 19, 38  
 g 1, 13      h 1, 2, 3, 6, 7, 14, 21, 42  
 i 1, 3, 5, 9, 15, 45  
 j 1, 2, 3, 4, 6, 9, 12, 18, 36
- 4 a 12, 24, 36      b 20, 40, 60  
 c 15, 30, 45      d 18, 36, 54  
 e 35, 70, 105
- 5 square numbers
- 6 13 is a prime number
- 7 2, 3, 5, 7, 11, 13, 17, 19
- 8  $1 + 3 + 5 + 7 + 9 = 25$   
 $1 + 3 + 5 + 7 + 9 + 11 = 36$   
 $1 + 3 + 5 + 7 + 9 + 11 + 13 = 49$   
 $1 + 3 + 5 + 7 + 9 + 11 + 13 + 15 = 64$

9

	Square number	Factor of 70
Even number	16	10
Multiple of 7	49	35

- 10 4761 ( $69^2$ )
- 11 4 packs of sausages, 5 packs of buns
- 12 24 seconds
- 13 30 seconds
- 14 12 minutes; Debbie: 4 and Fred: 3
- 15 a 12      b 9      c 6      d 13  
 e 15      f 14      g 16      h 10  
 i 18      j 17      k 8      l 21
- 16 b 21, 28, 36, 45, 55

**Exercise 5H**

- 1 a 2      b 5      c 7      d 1  
 e 9      f 10      g 8      h 3  
 i 6      j 4      k 11      l 12  
 m 20      n 30      o 13
- 2 a +5, -5      b +6, -6      c +10, -10  
 d +7, -7      e +8, -8      f +4, -4  
 g +3, -3      h +9, -9      i +1, -1  
 j +12, -12
- 3 a 81      b 40      c 100      d 14  
 e 36      f 15      g 49      h 12  
 i 25      j 21      k 121      l 16  
 m 64      n 17      o 441
- 4 a 24      b 31      c 45      d 40  
 e 67      f 101      g 3.6      h 6.5  
 i 13.9      j 22.2
- 5  $\sqrt{50}$ ,  $3^2$ ,  $\sqrt{90}$ ,  $4^2$
- 6 4 and 5
- 7  $\sqrt{324} = 18$
- 8 15 tiles
- 9 a 0.36      b 0.6      c  $\frac{3}{5}$   
 d i  $\frac{2}{3}$       ii  $\frac{5}{7}$       iii  $\frac{8}{9}$
- 10 Because  $8^2 = 64$  and  $9^2 = 81$  or  $\sqrt{64} = 8$  and  $\sqrt{81} = 9$ . 8.7 is between 8 and 9
- 11 a False      b True  
 c False      d True
- 12 a  $2^2 \times 3^4$       b  $2 \times 3^2 = 18$       c 22
- 13 a 16 900      b 3600

**Exercise 5I**

- 1 a 45      b 52      c 66
- 2 a 200      b 115      c 236
- 3 a 56      b 157      c 76      d 193
- 4 a 144      b 108
- 5 a 12.54      b 27.45
- 6 a 26.7      b 24.5      c 145.3      d 1.5
- 7 Sovereign is 102.48p per litre, Bridge is 102.73p per litre so Sovereign is cheaper.
- 8 Abby 1.247, Bobby 2.942, Col 5.333, Donna 6.538  
 Col is correct.
- 9  $31 \times 3600 \div 1610 = 69.31677 \approx 70$

10 a 167.552 b 196.48

11 a 2.77 b 6.00

12 a 497.952 b 110.978 625

13 a True b False

## Review questions

1 a 90 b 105

2 a Last digits: 4, 6, 4.  $4 \times 4 \times 4 \times 4 \times 4 = 1024$   
b 4

3 a 6, 12, 18, 24, 30 b 1, 2, 3, 4, 6, 12  
c 25 d 23, 29

4 13.69

5 53, 59

6 a 8 b 8 c 125

7 9<sup>th</sup> February

8 5 packets of pies and 2 packets of breadsticks

9 a  $2 \times 3 \times 3 \times 5$  b 5

10 630

11 a 15 b 3150

12 a 196 b  $35 \times 35$  must end in 5

13 a Clock A 3 am, clock B 3.15 am  
b Clock A 12 midday, clock B 1.00 pm

14 a 30 seconds b 50 seconds  
c 75 seconds d 150 seconds

15 a Newton's method = 7.746103896,  $\sqrt{60}$   
from calculator = 7.745966692 so the root  
is correct to 2 dp or 3dp if rounded.  
b Newton's method = 6.245 1612 9,  $\sqrt{39}$   
from calculator = 6.244 997 998 so  
the root is correct to 2 dp or 3 dp if  
rounded.

16  $x = y$

## Chapter 6 – Number: Approximations

### Exercise 6A

1 a 20 b 60 c 80 d 50  
e 100 f 20 g 90 h 70  
i 10 j 30 k 30 l 50  
m 80 n 50 o 90

2 a 200 b 600 c 800 d 500  
e 1000 f 100 g 600 h 400  
i 1000 j 1100 k 400 l 700  
m 800 n 1000 o 1100

3 a 1 b 2 c 1 d 1  
e 3 f 2 g 3 h 2  
i 1 j 1 k 3 l 2  
m 74 n 126 o 184

4 a 2000 b 6000 c 8000 d 5000  
e 10 000 f 1000 g 6000 h 3000  
i 9000 j 2000 k 3000 l 5000  
m 8000 n 5000 o 9000 p 4000  
q 7000 r 8000 s 1000 t 2000

5 a 230 b 570 c 720 d 520  
e 910 f 230 g 880 h 630  
i 110 j 300 k 380 l 630  
m 350 n 1010 o 1070

6 a True b False c True  
d True e True f False

7 Welcome to Swinton. Population 1400  
(to the nearest 100)

8 a Man Utd v Chelsea  
b QPR v Aston Villa  
c 20 000, 45 000, 75 000, 18 000, 30 000,  
44 000, 20 000, 36 000, 25 000, 35 000  
d 19 900, 44 600, 75 300, 18 000, 30 000,  
44 400, 20 300, 35 700, 24 700, 35 000

9 a 35 min b 55 min c 15 min  
d 50 min e 10 min f 15 min  
g 45 min h 35 min i 5 min  
j 0 min

10 a 375 b 350 to 374 inclusive

11 A number between 75 and 84 inclusive  
added to a number between 45 and 54  
inclusive with a total not equal to 130, for  
example  $79 + 49 = 128$

### Exercise 6B

1 a 4.8 b 3.8 c 2.2 d 8.3  
e 3.7 f 46.9 g 23.9 h 9.5  
i 0.1 j 0.6 k 65.0 l 51.0

2 a 5.78 b 2.36 c 0.98 d 33.09  
e 23.57 f 91.79 g 8.00 h 2.31  
i 6.00 j 3.51 k 96.51 l 0.01

3 a 4.6 b 0.08 c 45.716 d 94.85  
e 602.1 f 671.76 g 7.1 h 6.904  
i 13.78 j 0.1 k 4.002 l 60.0

- 4 a 9            b 9            c 7            d 3  
 e 8            f 8            g 6            h 4  
 i 2            j 47            k 154        l 343

5  $£1 + £7 + £4 + £1 = £13$

6 3, 3.46, 3.5

7 4.7275 or 4.7282

8 6.445 or 6.446 or 6.447 or 6.448 or 6.449

9  $8.75 \leq \text{value} < 8.85$

10 £8.45 to £8.49 inclusive

- 11 a  $6.45 \text{ cm} \leq \text{value} < 6.55 \text{ cm}$   
 b Smallest value of perimeter =  
 $2 \times (6.45 + 3.25) = 19.4 \text{ cm}$   
 Largest value of perimeter =  
 $2 \times (6.55 + 3.35) = 19.8 \text{ cm}$   
 Therefore  $19.4 \leq \text{perimeter} < 19.8$   
 c Smallest area =  $6.45 \times 3.25 = 20.96 \text{ cm}^2$

12 a 29.83 cm            b 32.991 cm

13 3.645

### Exercise 6C

- 1 a 50 000            b 60 000            c 30 000  
 d 90 000            e 90 000            f 50  
 g 90            h 30            i 100  
 j 200            k 0.5            l 0.3  
 m 0.006            n 0.05            o 0.0009  
 p 10            q 90            r 90  
 s 200            t 1000

- 2 a 56 000            b 27 000            c 80 000  
 d 31 000            e 14 000            f 5900  
 g 1100            h 850            i 110  
 j 640            k 1.7            l 4.1  
 m 2.7            n 8.0            o 42  
 p 0.80            q 0.46            r 0.066  
 s 1.0            t 0.0098

- 3 a 64 500            b 19 300            c 4.55  
 d 4.08            e 14.4            f 6.69  
 g 1.07            h 9430            i 10.4  
 j 9.68

4 a 65, 74            b 95, 149            c 950, 1499

5 Ayton: 750, 849; Beeville: 950, 1499;  
 Charlestown: 150 000, 249 999

6 a 64.4            b 63.5

7 15, 16 or 17

8 1, because there could be 450 then 449

9 Between 1945 and 1954

### Exercise 6D

- 1 a 35 000            b 15 000            c 960  
 d 12 000            e 1050            f 4000  
 g 5            h 20            i 1200

2 a £4000            b £3000            c £2000            d £800

3 a £15 000            b £20 000            c £15 000

4 £21 000

- 5 a 14            b 10            c 1.1            d  $\frac{3}{4}$   
 e 5            f  $\frac{2}{3}$             g 3            h 0.3  
 i 6            j 400            k 2            l 24

- 6 a 500            b 200            c 90  
 d 50            e 50            f 500

7 8

8 a 200            b 2800            c 10            d 1000

9 a 40            b 10            c £70

10 1000 or 1200

11 Yes, the approximation for both distances is 120 km

12 400

13 a 3 kg            b 200

14 a \$210 000 000            b \$42 000 000

15  $7.5 \leq N < 8.5$

16 a  $9.5 \leq L < 10.5$             b  $4.5 \leq W < 5.5$   
 c  $28.0 \leq P < 32.0$             d  $42.75 \leq A < 57.75$

### Review questions

1 a 6000            b 790

2 a 3200            b 5000

3 a i 36 000 000            ii 3 600  
 b i 1000            ii 100

4 a  $259 \approx 260 \text{ m}$             b  $1340 - 1090 = 250$   
 c No, one value rounds up, the other down

5 a 7650            b 7749

6 a 2450            b 2549            c i 2475            ii 2524

7 a 2            b 3            c 3            d 4

8 a 2            b 1            c 2            d 3  
 e 3            f 4            g 3            h 1

9 a 67.5            b 31            c 3000            d 4.0  
 e 20            f 12.4            g 13.57            h 29.4

- 10 No, as there could be 44 people on the bus and 24 at the bus stop. That makes 68 people altogether and the bus can only have  $52 + 10 = 62$  passengers maximum.
- 11 Width wise: Yes, as error interval of the envelope is  $8.5 \leq \text{width} < 9.5$  and for the postcard the error interval is  $7.75 \leq \text{width} < 8.25$
- Length wise: No, as error interval of the envelope is  $12.5 \leq \text{length} < 13.5$  and for the postcard the error interval is  $12.25 \leq \text{length} < 12.75$ , so largest postcard is larger than the smallest envelope.
- 12 1.850 or 1.851 or 1.852 or 1.853 or 1.854
- 13  $235 \leq \text{number} < 245$
- 14 a  $49.5 \leq \text{side} < 50.5$   
 b No, as the error interval for the peg is  $48 \leq \text{diameter} < 50$
- 15 a  $10.5 \leq \text{perimeter} < 13.5$   
 b  $4.375 \leq \text{area} < 7.875$

## Chapter 7 – Number: Decimals and fractions

### Exercise 7A

- 1 a 0.48    b 2.92    c 1.12    d 0.12  
 e 0.028    f 0.09    g 0.192    h 2.91
- 2 a 8    b 9    c 4    d 8.7  
 e 7.8    f 9.7    g 12.1    h 9.7
- 3 Yes, with 25p left over
- 4 Alistair 0.90 Shane 70 Charlotte 66
- 5 27 (three 0.7 m lengths from each 2.4 m length)
- 6 a i  $5 \times 1 = 5$     ii 3.36, 1.64  
 b i  $2 \times .5 = 1$     ii 1.44, 0.44  
 c i  $15 \times 1 = 15$     ii 13.77, 1.23  
 d i  $20 \times 1 = 20$     ii 16.08, 3.92
- 7 a 572    b i 5.72    ii 1.43    iii 22.88
- 8 a 29    b 24    c 0.12    d 0.08
- 9 a  $33.25 \leq ab < 47.25$   
 b Because it is the highest value of a divided by the lowest value of b  
 c 2.11

### Exercise 7B

- 1 a 0.5    b  $0.\dot{3}$     c 0.25    d 0.2  
 e  $0.1\dot{6}$     f  $0.\dot{1}4285\dot{7}$     g 0.125    h  $0.\dot{1}$   
 i 0.1    j  $0.\dot{0}76\ 92\dot{3}$
- 2 a i 0.571428571428571428571428 ...  
 ii 0.714285714285714285714285 ...  
 iii 0.857142857142857142857142 ...
- b The recurring digits are all in the same sequence but they start in a different place each time.
- 3 a i 0.4    ii 0.2    iii 0.1  
 b Halve the answers to get  $1 \div 20 = 0.05$  and  $1 \div 40 = 0.025$
- 4  $0.\dot{1}$ ,  $0.\dot{2}$ ,  $0.\dot{3}$ ,  $0.\dot{4}$ ,  $0.\dot{5}$ ,  $0.\dot{6}$ ,  $0.\dot{7}$ ,  $0.\dot{8}$   
 The recurring digit is the numerator of the fraction.
- 5  $0.\dot{0}9$ ,  $0.\dot{1}8$ ,  $0.\dot{2}7$ ,  $0.\dot{3}6$ ,  $0.\dot{4}5$ ,  $0.\dot{5}4$ ,  $0.\dot{6}3$ ,  $0.\dot{7}2$ ,  $0.\dot{8}1$ ,  $0.\dot{9}0$   
 The recurring digits follow the nine times table.
- 6  $\frac{9}{22} = 0.4\dot{0}9$ ,  $\frac{3}{7} = 0.428\ 57\dot{1}$ ,  
 $\frac{16}{37} = 0.432$ ,  $\frac{4}{9} = 0.\dot{4}$ ,  
 $\frac{5}{11} = 0.\dot{4}5$ ,  $\frac{6}{13} = 0.46153\dot{8}$ ,
- 7  $\frac{7}{24} = \frac{35}{120}$ ,  $\frac{3}{10} = \frac{36}{120}$ ,  $\frac{19}{60} = \frac{38}{120}$ ,  
 $\frac{2}{5} = \frac{48}{120}$ ,  $\frac{5}{12} = \frac{50}{120}$
- 8 a  $\frac{1}{8}$     b  $\frac{17}{50}$     c  $\frac{29}{40}$     d  $\frac{5}{16}$   
 e  $\frac{89}{100}$     f  $\frac{1}{20}$     g  $2\frac{7}{20}$     h  $\frac{7}{32}$
- 9 a 0.08 $\dot{3}$     b 0.0625    c 0.05  
 d 0.04    e 0.02
- 10 a  $\frac{4}{3} = 1\frac{1}{3}$     b  $\frac{6}{5} = 1\frac{1}{5}$     c  $\frac{5}{2} = 2\frac{1}{2}$   
 d  $\frac{10}{7} = 1\frac{3}{7}$     e  $\frac{20}{11} = 1\frac{9}{11}$     f  $\frac{15}{4} = 3\frac{3}{4}$
- 11 a 0.75,  $1.\dot{3}$     b  $0.8\dot{3}$ , 1.2    c 0.4, 2.5  
 d 0.7,  $1.428\ 57\dot{1}$     e 0.55,  $1.8\dot{1}$     f  $0.2\dot{6}$ , 3.75  
 b No, look at    c above for example.
- 12 The answer is always 1.

13  $1 \div 0$  is infinite so there is no finite answer.

14 a 10      b 2  
c The reciprocal of a reciprocal is always the original number.

15 The reciprocal of  $x$  is greater than the reciprocal of  $y$ .  
For example, reciprocal of 2 is 0.5, reciprocal of 10 is 0.1.

### Exercise 7C

1 a  $\frac{1}{3}$       b  $\frac{1}{5}$       c  $\frac{2}{5}$       d  $\frac{5}{24}$   
e  $\frac{2}{5}$       f  $\frac{1}{6}$       g  $\frac{2}{7}$       h  $\frac{1}{3}$

2  $\frac{3}{5}$

3  $\frac{2}{5}$

4  $\frac{2}{3}$

5  $\frac{1}{5}$

6  $\frac{1}{2}$

7 Jon saves  $\frac{30}{90} = \frac{1}{3}$   
Matt saves,  $\frac{35}{100} = \frac{7}{20}$  which is greater than  $\frac{1}{3}$ , so Matt saves the greater proportion of his earnings.

8  $\frac{13}{20} = \frac{65}{100}$  and  $\frac{16}{25} = \frac{64}{100}$  so 13 out of 20 is the better mark.

9  $\frac{3}{8}$

10  $\frac{11}{24}$

11 26 out of 63 because 13 out of 32 is the same as 26 out of 64.

12 The price is the same at both shops: £432

### Exercise 7D

1 a  $\frac{8}{15}$       b  $\frac{7}{12}$       c  $\frac{11}{12}$   
d  $\frac{1}{10}$       e  $\frac{1}{8}$       f  $\frac{1}{12}$

2 Three-quarters of 68

3 a  $4\frac{47}{60}$       b  $\frac{41}{72}$       c  $1\frac{109}{120}$       d  $1\frac{23}{30}$

4 a  $\frac{1}{6}$       b 30

5 No, one eighth is left, which is 12.5 cl, so enough for one cup but not two cups.

6 He has added the numerators and added the denominators instead of using a common denominator. Correct answer is  $3\frac{7}{12}$ .

7 Possible answer: The denominators are 4 and 5. I first find a common denominator. The lowest common denominator is 20 because 4 and 5 are both factors of 20. So I am changing the fractions to twentieths. One-quarter is the same as five-twentieths (multiplying numerator and denominator by 5). Two-fifths is the same as eight-twentieths (multiplying numerator and denominator by 4). Five-twentieths plus eight-twentieths = thirteen-twentieths.

8  $\frac{11}{20}$  of 900 = 495,  $\frac{2}{11}$  of 495 = 90 left-handed boys.  $900 - 495 = 405$  girls.  $\frac{2}{9}$  of 405 = 90 left-handed girls. 180 left-handed students altogether so 180 out of 900 =  $\frac{1}{5}$ .

9  $\frac{1}{5} + \frac{3}{8} = \frac{23}{40}$ , so  $\frac{17}{40}$  of the counters are yellow.  $\frac{17}{40}$  of 600 = 255

10 a because  $\frac{27}{40} + \frac{2}{5} = 1\frac{3}{40}$  which is greater than 1.

b  $\frac{2}{5}$  of 200 = 80.  $\frac{5}{8}$  of 80 = 50 women at least 40.  $\frac{27}{40}$  of 200 = 135 members at least 40.  $135 - 50 = 85$  men at least 40.  $\frac{3}{5}$  of 200 = 120, so  $120 - 85 = 35$  men under 40.

### Exercise 7E

1 a  $\frac{1}{6}$       b  $\frac{3}{8}$       c  $\frac{7}{20}$       d  $\frac{3}{5}$   
e  $\frac{5}{12}$       f  $2\frac{11}{12}$       g  $3\frac{9}{10}$       h  $3\frac{1}{3}$

2 a  $\frac{3}{4}$       b  $1\frac{1}{15}$       c 5

d  $\frac{4}{9}$       e  $1\frac{3}{5}$

3  $\frac{3}{8}$

- 4  $\frac{1}{8}$   
 5 40  
 6  $\frac{2}{5}$  of  $6\frac{1}{2}$   
 7 £10.40

**Exercise 7F**

- 1 a  $1\frac{11}{20}$       b  $1\frac{1}{4}$       c  $1\frac{63}{80}$   
 d  $\frac{11}{30}$       e  $\frac{61}{80}$       f  $\frac{167}{240}$
- 2 a  $12\frac{1}{4}$  miles      b  $3\frac{1}{4}$  miles
- 3 a  $6\frac{11}{20}$       b  $8\frac{8}{15}$       c  $11\frac{63}{80}$   
 d  $3\frac{11}{30}$       e  $7\frac{61}{80}$       f  $1\frac{277}{396}$
- 4 a  $-\frac{77}{1591}$       b The answer is negative
- 5  $\frac{5}{12}$  (anticlockwise) or  $\frac{7}{12}$  (clockwise)
- 6 a  $\frac{3}{5}$       b  $\frac{27}{128}$       c  $5\frac{2}{5}$       d  $5\frac{1}{7}$   
 e  $3\frac{9}{32}$       f  $\frac{11}{18}$
- 7 a  $8\frac{11}{20}$       b  $65\frac{91}{100}$       c  $52\frac{59}{160}$       d  $2\frac{17}{185}$   
 e  $2\frac{22}{103}$       f  $7\frac{881}{4512}$
- 8 3 rolls
- 9  $22 \div (2 \times \frac{22}{7}) = \frac{7}{2}$ ,  $\frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} = 38\frac{1}{2}$  cm<sup>2</sup>

**Review questions**

- 1 0.3 < decimal < 0.33... for example 0.32
- 2 17
- 3 a 0.4      b  $\frac{9}{20}$
- 4  $\frac{22}{75}$
- 5 a 0.5      b 0.3
- 6  $28 \div 40 = 0.7$ ;  $33 \div 50 = 0.66$ . Zac did better in English
- 7  $\frac{4}{5}$
- 8  $900 \times 12 = 10\ 800$  miles per year.  $10\ 800 \div 45 = 240$  gallons.  $240 \times 4.55 = 1092$  litres.  $1092 \times 1.4 = £1528.80$  per year.

- 9 a 5      b  $\frac{3}{2}$       c  $\frac{7}{10}$       d  $\frac{1}{8}$
- 10 Starbuys  $10 \times 0.8 = £8$ . Shoprite  $12 \times 0.8 = £9.60$ . 20% off = £7.68, so Shoprite is best value.

- 11 a 1672      b i 167.2      ii 83.6      iii 16.72

- 12 a 3.2      b 360      c 0.32

- 13 a  $\frac{7}{20}$       b  $\frac{2}{5}$       c  $1\frac{1}{2}$

- 14 40.96

- 15  $2\frac{1}{24}$

- 16  $18\frac{11}{12}$  cm

- 17  $18\frac{5}{12}$

- 18 a 0.5 (T), 0.6 (R), 0.75 (T), 0.8 (T), 0.83 (R), 0.857142 (R), 0.875 (T), 0.8 (R), 0.9 (T)  
 b 2,  $2 \times 2$ , 5,  $2 \times 2 \times 2$ ,  $2 \times 5$   
 c If the prime factors of the denominator of a fraction in its simplest form are only 2 and/or 5 its decimal will terminate.

19

	Blue	Red	Total
Squares	136	74	210
Triangles	95	95	190
Total	231	169	400

**Chapter 8 – Algebra: Linear graphs**

**Exercise 8A**

- 1 a A(1, 2), B(3, 0), C(0, 1), D(-2, 4), E(-3, 2), F(-2, 0), G(-4, -1), H(-3, -3), I(1, -3), J(4, -2)  
 b i (2, 1)      ii (-1, -3)      iii (1, 1)
- 2 a  $x = -3$ ,  $x = 2$ ,  $y = 3$ ,  $y = -4$   
 b i  $x = -\frac{1}{2}$       ii  $y = -\frac{1}{2}$
- 3 Values of y: 2, 3, 4, 5, 6
- 4 Values of y: -2, 0, 2, 4, 6
- 5 Values of y: 1, 2, 3, 4, 5
- 6 Values of y: -4, -3, -2, -1, 0
- 7 a Values of y: 0, 4, 8, 12, 16 and 6, 8, 10, 12, 14  
 b (6, 12)
- 8 Points could be (0, -1), (1, 4), (2, 9), (3, 14), (4, 19), (5, 24), etc



- b** £25.00, £28.50, £32.00, £35.50, £39.00, £42.50;  
 (£20.00), (£23.50), (£27.00), (£30.50), (£34.00), (£37.50);  
 (£15.00), (£18.50), (£22.00), (£25.50), (£29.00), (£32.50);  
 (£10.00), (£13.50), (£17.00), (£20.50), (£24.00), (£27.50);  
 (£5.00), (£8.50), (£12.00), (£15.50), (£19.00), (£22.50);  
 (£0.00), (£3.50), (£7.00), (£10.50), (£14.00), (£17.50)

**c** Yes, they had 3 rolls and 4 baguettes.



**b**  $y = 3x + 1$





**c** Graph from (0, 1) to (5, 16)

**d** Read from 13 on the  $y$ -axis across to the graph and down to the  $x$ -axis. This should give a value of 4.


**11** 0, 8, 16, 24, 32, 40, 48

### Exercise 8B


**1**




**2**



**3**




**4 a**




**b** (2, 7)

**5 a**



**b**




**c** Lines in part **a** intersect at (6, 1). Lines in part **b** don't intersect because they are parallel.

6 a Line isn't straight


b

x	-3	-2	-1	0	1	2	3
y	-5	-3	-1	1	3	5	7




b Ian, as Ian only charges £85, whilst Joan charges £90 for a 2-hour job

8 Jada's




b 4.5 units squared



Exercise 8C

- 1 a 2      b  $\frac{1}{3}$       c -3      d 1  
 e -2      f  $-\frac{1}{3}$       g 5      h -5  
 i  $\frac{1}{5}$       j  $-\frac{3}{4}$



- 3 a Both answers are correct  
 b generally the bigger the triangle the more accurate the answer, so Brianna
- 4 a Slide won't be steep enough for children to slide down.  
 b If it is too steep children might slide down too quickly.  
 c A, B, E and F satisfy the safety regulations; C and D do not

5 a 0.5      b 0.4      c 0.2      d 0.1      e 0

6 a 1.4      b 2      c 3.3      d 10      e  $\infty$


7 Raisa has misread the scales. The second line has four times the gradient (2.4) of the first (0.6).

8 4 horizontal 6 vertical; 5 horizontal 8 vertical; 3 horizontal 5 vertical; 6 horizontal 11 vertical; 1 horizontal 2 vertical; 2 horizontal 5 vertical


9 0, 2, -1,  $\frac{1}{2}$ ,  $-\frac{3}{2}$

Exercise 8D


1 a, b, c, d




e, f, g, h




l, j, k, l




e, f, g, h



2 a



l, j, k, l



b (-6, 0)

3 a  $y = 3x - 2$


b They intercept the y-axis at the same point (0, -2).

c (-1, -4)


4 a -2      b  $\frac{1}{2}$

**Exercise 8E**

1 a, b, c, d



2 a



b (2, 2)

3 a Intersect at (6, 0)

b Intersect at (0, -3)

c Parallel

d  $-2x + 9y = 18$

4 a vi

b iii

c v

d ii

e i

f iv

5 a i  $x = 3$

ii  $x - y = 4$

iii  $y = -3$

iv  $x + y = -4$

v  $x = -3$

vi  $y = x + 4$

b i -3

ii  $\frac{1}{3}$

iii  $-\frac{1}{3}$

- 6 Cover-up method for  $2x + y = 10$  and gradient-intercept method for  $y = 1 - 2x$ , but actually the lines are the same

**Exercise 8F**

- 1 a  $y = \frac{4}{3}x - 2$  or  $3y = 4x - 6$   
 b  $y = 2x$       c  $2y = x + 6$
- 2 a i  $y = 2x + 1$ ,  $y = -2x + 1$   
 ii Reflection in  $y$ -axis (and  $y = 1$ )  
 iii Different sign  
 b i  $5y = 2x - 5$ ,  $5y = -2x - 5$   
 ii Reflection in  $y$ -axis (and  $y = -1$ )  
 iii Different sign  
 c i  $y = x + 1$ ,  $y = -x + 1$   
 ii Reflection in  $y$ -axis (and  $y = 1$ )  
 iii Different sign
- 3 a  $x$ -coordinates go from  $2 \rightarrow 1 \rightarrow 0$  and  $y$ -coordinates go from  $5 \rightarrow 3 \rightarrow 1$ .  
 b  $x$ -step between the points is 1 and  $y$ -step is 2.  
 c  $y = 3x + 2$
- 4 a  $y = -x + 1$   
 b  $5y = -2x - 5$   
 c  $y = -\frac{3}{2}x - 3$  or  $2y = -3x - 6$
- 5 a i  $y = \frac{1}{2}x$ ,  $y = 2x$   
 ii Reflection in  $x = y$   
 iii Reciprocal of each other  
 b i  $y = 2$ ,  $x = 2$       ii Reflection in  $x = y$   
 iii Reciprocal of each other (reciprocal of zero is infinity)
- 6 All of the lines except  $y = \frac{1}{4}x + 9$
- 7 a  $y = -3x + 5$       b  $y = 2x - 4$   
 c  $y = 8x - 3$       d  $y = 25 - 2x$   
 e  $y = \frac{2}{3}x - 1$
- 8  $5x + 6y = 30$
- 9 Chris is correct. The equation of the line is  $y = \frac{1}{2}x + 2$  and  $(12, 8)$  satisfies the rule
- 10 a i  $x + y = 100$       ii  $k = 1$   
 b i  $x = 46$       ii  $k = 46$   
 c i  $y = 2x + 1$       ii  $k = 60$   
 d i  $y = x + 19$       ii  $k = -17$

**Exercise 8G**

- 1 A and B are parallel because they have the same gradient ( $m = 3$ )

- 2  $x = 8$  and  $x = -2$   
 $y = 3 - x$  and  $y = 10 - x$   
 $y = 3x - 5$  and  $y = 3x + 9$   
 $y = -\frac{1}{8}x$  and  $y = -\frac{1}{8}x + 6$   
 $2x + y = 9$  and  $2x + y = 13$   
 $y = 15 - 4x$  and  $y = 7 - 4x$
- 3 a  $y = \frac{1}{2}x - 2$   
 b  $y = -x + 3$   
 c  $y = 3x - 8$   
 d  $y = -\frac{1}{3}x + 10$
- 4 a  $y = 2x + 4$       b  $y = 17 - 3x$   
 c  $y = 3x - 1$

**Exercise 8H**




- 1 a i  $8\frac{1}{4}$  kg      ii  $2\frac{1}{4}$  kg      iii 9 lb      iv 22 lb  
 b 2.2 lb  
 c Read off the value for 12 lb (5.4 kg) and multiply this by 4 (21.6 kg)  
 d  $m = \frac{5}{11}$   
 e  $\frac{5}{11}$  kilograms per pound
- 2 a i £100      ii £325  
 b i 500      ii 250
- 3 a i 95 °F      ii 68 °F      iii 10 °C      iv 32 °C  
 b 32 °F  
 c  $m = \frac{5}{9}$ ,  $\frac{5}{9}$  °C per °F
- 4 a Check student's graph  
 b 2:15 pm  
 c  $m = 48$ , 48 cups of tea per hour
- 5 a Check student's graph, £50
- 6 a Anya: CabCo £8.50, YellaCabs £8.40, so YellaCabs is best  
 Bettina: CabCo £11.50, YellaCabs £11.60, so CabCo is best  
 Calista: CabCo £10, YellaCabs £10, so either  
 b If they shared a cab, the shortest distance is 16 km, which would cost £14.50 with CabCo and £14.80 with Yellacabs.
- 7 a \$642.86  
 b i \$7500      ii £3666.67

**Exercise 8I**

- 1 (4, 1)  
 2 (2, 3)

- 3 (3, 10)  
 4 (-2, 6)  
 5 (-6, -9)  
 6 (1, -1)  
 7 (2, 6)  
 8 (2, 8)  
 9  $x + 2y = 9.5$ ,  $2x + y = 8.5$   
 Graphs intersect at (2.5, 3.5), so a bun costs £2.50 and a cake costs £3.50.  
 10 a P and R  
 b R and S  
 c P and Q  
 d Q and S  
 11 (0, 0), (-3, 3), (-3, -3), (-3, 2), (-2, 2), (2, 2)  
 12 a no solutions  
 b infinite solutions  
 c one solution  
 d a Parallel b Same line c Not parallel

## Review questions



- 4 a  $\frac{1}{10}$   
 b 24.5 cm  
 c 0.1 cm or 1 mm  
 d  $l = 24.5 + \frac{W}{10}$  or Length =  $24.5 + 1 \text{ mm/kg}$   
 5 A Q, B T, C P, D R, E S  
 6 a (5, 5)  
 b (1, 5)  
 c (3, 16)  
 7 72 square units  
 8 a 2 b  $y = 2x + 2$   
 c  $y = 2x - 5$   
 9 a  $x + y = 50$  (or  $y = 50 - x$ )  
 b -3  
 c  $x + y = 40$  (or  $y = 40 - x$ )

## Chapter 9 – Algebra: Expressions and formulae

### Exercise 9A

- 1 a  $x + 2$  b  $x - 6$  c  $hj$   
 d  $y \div t$  or  $\frac{y}{t}$  e  $a^2$  f  $g^2$   
 2 a Dad:  $(72 + x)$  years, me:  $(T + x)$  years b 31  
 3 For example,  $2 \times 6m$ ,  $1 \times 12m$ ,  $6m + 6m$ , etc  
 4 Any values picked for  $l$  and  $w$  and substituted into the formulae to give the same answers  
 5 13 years  
 6 15p  
 7 a expression, c identity, b formula, d equation

### Exercise 9B

- 1 a 3 b 11 c 43  
 2 a 9 b 5 c -1  
 3 a £4.00 b 13 km  
 c Yes, the journey will cost £5.00  
 4 a 1.4 b 1.4 c -0.4  
 5 a 2.5 b -20 c 2.5  
 6 a  $\frac{150}{n}$  b £925  
 7 a  $2 \times 8 + 6 \times 11 - 3 \times 2 = 76$   
 b  $5 \times 2 - 2 \times 11 + 3 \times 8 = 12$

8 a One odd, one even value, different from each other.

b Any valid combination, e.g.  $x = 1, y = 2$

9 a i Odd ii Odd iii Even iv Odd

b Any valid expression such as  $xy + z$

10 a £20 b -£40 c 40 miles

11 a First term is cost of petrol, each mile is a tenth of £1.20. Second term is the hire cost divided by the miles.

b 32 p per mile

### Exercise 9C

1 a  $6 + 2m$  b  $10 + 5l$  c  $12 - 3y$   
 d  $20 + 8k$  e  $6 - 12f$  f  $10 - 6w$   
 g  $10k + 15m$  h  $12d - 8n$  i  $t^2 + 3t$   
 j  $k^2 - 3k$  k  $4t^2 - 4t$  l  $8k - 2k^2$   
 m  $8g^2 + 20g$  n  $15h^2 - 10h$  o  $y^3 + 5y$   
 p  $h^4 + 7h$  q  $k^3 - 5k$  r  $3t^3 + 12t$   
 s  $15d^3 - 3d^4$  t  $6w^3 + 3tw$  u  $15a^3 - 10ab$   
 v  $12p^4 - 15mp$  w  $12h^3 + 8h^2g$  x  $8m^3 + 2m^4$

2 a  $5(t - 1)$  and  $5t - 5$   
 b Yes, as  $5(t - 1)$  when  $t = 4.50$  is  $5 \times 3.50 = £17.50$

3 a He has worked out  $3 \times 5$  as 8 instead of 15 and he has not multiplied the second term by 3.

b  $15x - 12$

4 a  $3(2y + 3)$   
 b  $2(6z + 4)$  or  $4(3z + 2)$

### Exercise 9D

1 a  $7t$  b  $3y$  c  $9d$  d  $3e$   
 e  $3p$  f  $2t$  g  $5t^2$  h  $5ab$   
 i  $3a^2d$

2 a  $22 + 5t$  b  $21 + 19k$  c  $22 + 2f$   
 d  $14 + 3g$

3 a  $2 + 2h$  b  $9g + 5$  c  $17k + 16$   
 d  $6e + 20$

4 a  $4m + 3p + 2mp$  b  $3k + 4h + 5hk$   
 c  $12r + 24p - 13pr$  d  $19km + 20k - 6m$

5 a  $9t^2 + 13t$  b  $13y^2 + 5y$   
 c  $10e^2 - 6e$  d  $14k^2 - 3kp$

6 a  $17ab + 12ac + 6bc$  b  $18wy + 6ty - 8tw$   
 c  $14mn - 15mp - 6np$  d  $8r^3 - 6r^2$

7 a  $5(f + 2s) + 2(2f + 3s) = 9f + 16s$   
 b i £270f ii £480s  
 c  $£42\,450 - £30\,000 = £12\,450$

8 For  $x$ -coefficients, 3 and 1 or 1 and 4; for  $y$ -coefficients, 5 and 1 or 3 and 4 or 1 and 7

9  $5(3x + 2) - 3(2x - 1) = 9x + 13$

### Exercise 9E

1 a  $6(m + 2t)$  b  $3(3t + p)$  c  $4(2m + 3k)$   
 d  $4(r + 2t)$  e  $m(n + 3)$  f  $g(5g + 3)$   
 g  $2(2w - 3t)$  h  $y(3y + 2)$  i  $t(4t - 3)$   
 j  $3m(m - p)$  k  $3p(2p + 3t)$  l  $2p(4t + 3m)$   
 m  $4b(2a - c)$  n  $5bc(b - 2)$  o  $2b(4ac + 3de)$   
 p  $2(2a^2 + 3a + 4)$  q  $3b(2a + 3c + d)$   
 r  $t(5t + 4 + a)$  s  $3mt(2t - 1 + 3m)$   
 t  $2ab(4b + 1 - 2a)$  u  $5pt(2t + 3 + p)$

2 a Mary has taken out a common factor.  
 b Because the bracket adds up to £10 c £30

3 a, d, f and h do not factorise  
 b  $m(5 + 2p)$  c  $t(t - 7)$   
 e  $2m(2m - 3p)$  g  $a(4a - 5b)$   
 i  $b(5a - 3bc)$

4 a Bernice  
 b Aidan has not taken out the largest possible common factor. Craig has taken  $m$  out of both terms but there isn't an  $m$  in the second term.

5 There are no common factors.

6 Perimeter =  $2x + 8 + x + 5 + 5x + 4 + 9x - 3 + 10 - x = 16x + 24 = 8(2x + 3)$

7  $\frac{4x^2 - 12x}{2x - 6} = \frac{4x(x - 3)}{2(x - 3)}$

### Exercise 9F

1 a  $x^2 + 5x + 6$  b  $t^2 + 7t + 12$   
 c  $w^2 + 4w + 3$  d  $m^2 + 6m + 5$

2 a  $p^2 + 3p - 70$  b  $u^2 - 12u + 32$   
 c  $k^2 + 2k - 15$  d  $z^2 - 12z + 27$

3 a should be 35 on the end  
 b should be -80 c should be  $-10x$   
 d should be 12y e should be  $-9z$

### Exercise 9G

1 a  $k^2 + 8k + 15$  b  $a^2 + 5a + 4$   
 c  $x^2 + 2x - 8$  d  $t^2 + 2t - 15$   
 e  $w^2 + 2w - 3$  f  $f^2 - f - 6$

2 a  $r^2 - 10r + 16$  b  $s^2 - 17s + 70$   
 c  $d^2 - 17d + 16$  d  $m^2 - 9m + 18$   
 e  $q^2 - 20q + 99$  f  $y^2 - 13y + 40$

3 a  $20a$  b  $3b$  c 200  
 d  $-11d$  e  $12e, 28$

**Exercise 9H**

- 1 a  $g^2 - 3g - 4$       b  $y^2 + y - 12$   
 c  $x^2 + x - 12$       d  $p^2 - p - 2$   
 e  $k^2 - 2k - 8$       f  $y^2 + 3y - 10$   
 g  $a^2 + 2a - 3$
- 2 a  $x^2 - 9$     b  $t^2 - 25$     c  $m^2 - 16$   
 d  $t^2 - 4$     e  $y^2 - 64$     f  $p^2 - 1$   
 g  $25 - x^2$     h  $49 - g^2$     i  $x^2 - 36$
- 3  $(x + 2)$  and  $(x + 3)$
- 4 a B:  $1 \times (x - 2)$  C:  $1 \times 2$  D:  $2 \times (x - 1)$   
 b  $(x - 2) + 2 + 2(x - 1) = 3x - 2$   
 c Area A =  $(x - 1)(x - 2)$  = area of square minus areas (B + C + D) =  $x^2 - (3x - 2) = x^2 - 3x + 2$
- 5 a  $x^2 + 2x + 1$     b  $x^2 - 2x + 1$     c  $x^2 - 1$

**Exercise 9I**

- 1 a  $6x^2 + 11x + 3$       b  $12y^2 + 17y + 6$   
 c  $6t^2 + 17t + 5$       d  $8t^2 + 2t - 3$   
 e  $10m^2 - 11m - 6$     f  $12k^2 - 11k - 15$   
 g  $6p^2 + 11p - 10$     h  $10w^2 + 19w + 6$   
 i  $6a^2 - 7a - 3$       j  $8r^2 - 10r + 3$   
 k  $15g^2 - 16g + 4$     l  $12d^2 + 5d - 2$   
 m  $8p^2 + 26p + 15$     n  $6t^2 + 7t + 2$   
 o  $6p^2 + 11p + 4$       p  $6 - 7t - 10t^2$   
 q  $12 + n - 6n^2$       r  $6f^2 - 5f - 6$   
 s  $12 + 7q - 10q^2$     t  $3 - 7p - 6p^2$   
 u  $4 + 10t - 6t^2$
- 2 a  $(3x - 2)(2x + 1) = 6x^2 - x - 2$ ,  
 $(2x - 1)(2x - 1) = 4x^2 - 4x + 1$ ,  
 $(6x - 3)(x + 1) = 6x^2 + 3x - 3$ ,  
 $(4x + 1)(x - 1) = 4x^2 - 3x - 1$   
 $(3x + 2)(2x + 1) = 6x^2 + 7x + 2$
- 3 a  $4x^2 - 1$     b  $9t^2 - 4$     c  $25y^2 - 9$   
 d  $16m^2 - 9$     e  $4k^2 - 9$     f  $16h^2 - 1$   
 g  $4 - 9x^2$     h  $25 - 4t^2$     i  $36 - 25y^2$   
 j  $a^2 - b^2$     k  $9t^2 - k^2$     l  $4m^2 - 9p^2$   
 m  $25k^2 - g^2$     n  $a^2b^2 - c^2d^2$     o  $a^4 - b^4$
- 4 a  $a^2 - b^2$   
 b Dimensions:  $a + b$  by  $a - b$ ; Area:  $a^2 - b^2$   
 c Areas are the same, so  $a^2 - b^2 = (a + b) \times (a - b)$
- 5 First shaded area is  $(2k)^2 - 1^2 = 4k^2 - 1$   
 Second shaded area is  $(2k + 1)(2k - 1) = 4k^2 - 1$

**Exercise 9J**

- 1 a  $x^2 + 10x + 25$       b  $m^2 + 8m + 16$   
 c  $t^2 + 12t + 36$       d  $p^2 + 6p + 9$   
 e  $m^2 - 6m + 9$       f  $t^2 - 10t + 25$   
 g  $m^2 - 8m + 16$       h  $k^2 - 14k + 49$

- 2 a  $9x^2 + 6x + 1$       b  $16t^2 + 24t + 9$   
 c  $25y^2 + 20y + 4$     d  $4m^2 + 12m + 9$   
 e  $16t^2 - 24t + 9$       f  $9x^2 - 12x + 4$   
 g  $25t^2 - 20t + 4$       h  $25r^2 - 60r + 36$   
 i  $x^2 + 2xy + y^2$       j  $m^2 - 2mn + n^2$   
 k  $4t^2 + 4ty + y^2$       l  $m^2 - 6mn + 9n^2$   
 m  $x^2 + 4x$             n  $x^2 - 10x$   
 o  $x^2 + 12x$             p  $x^2 - 4x$
- 3 a Bernice has just squared the first term and the second term. She hasn't written down the brackets twice.  
 b Pete has written down the brackets twice but has worked out  $(3x)^2$  as  $3x^2$  and not  $9x^2$ .  
 c  $9x^2 + 6x + 1$
- 4 Whole square is  $(2x)^2 = 4x^2$ .  
 Three areas are  $2x - 1$ ,  $2x - 1$  and 1.  
 $4x^2 - (2x - 1 + 2x - 1 + 1) = 4x^2 - (4x - 1) = 4x^2 - 4x + 1$

**Exercise 9K**

- 1 a  $(x + 2)(x + 3)$       b  $(t + 1)(t + 4)$   
 c  $(m + 2)(m + 5)$       d  $(k + 4)(k + 6)$
- 2 a  $(p + 2)(p + 12)$     b  $(r + 3)(r + 6)$   
 c  $(w + 2)(w + 9)$       d  $(x + 3)(x + 4)$
- 3 a  $(a + 2)(a + 6)$       b  $(k + 3)(k + 7)$   
 c  $(f + 21)(f + 1)$       d  $(b + 8)(b + 12)$
- 4 a  $(t - 2)(t - 3)$       b  $(d - 4)(d - 1)$   
 c  $(g - 2)(g - 5)$       d  $(x - 3)(x - 12)$
- 5 a  $(c - 2)(c - 16)$     b  $(t - 4)(t - 9)$   
 c  $(y - 4)(y - 12)$     d  $(j - 6)(j - 8)$
- 6 a  $(p - 3)(p - 5)$       b  $(y + 6)(y - 1)$   
 c  $(t + 4)(t - 2)$       d  $(x + 5)(x - 2)$
- 7 a  $(m + 2)(m - 6)$     b  $(r + 1)(r - 7)$   
 c  $(n + 3)(n - 6)$       d  $(m + 4)(m - 11)$
- 8 a  $(w + 4)(w - 6)$     b  $(t + 9)(t - 10)$   
 c  $(h + 8)(h - 9)$       d  $(t + 7)(t - 9)$
- 9 a  $(d + 1)^2$             b  $(y + 10)^2$   
 c  $(t - 4)^2$             d  $(m - 9)^2$
- 10 a  $(x - 12)^2$           b  $(d + 3)(d - 4)$   
 c  $(t + 4)(t - 5)$       d  $(q + 7)(q - 8)$
- 11  $(x + 2)(x + 3)$ , giving areas of  $2x$  and  $3x$ , or  $(x + 1)(x + 6)$ , giving areas of  $x$  and  $6x$ .
- 12 a  $x^2 + (a + b)x + ab$   
 b i  $p + q = 7$   
 ii  $pq = 12$   
 c 7 can only be  $1 \times 7$  and  $1 + 7 \neq 12$

### Exercise 9L

- 1 **a**  $(x+3)(x-3)$       **b**  $(t+5)(t-5)$   
**c**  $(m+4)(m-4)$       **d**  $(3+x)(3-x)$   
**e**  $(7+t)(7-t)$       **f**  $(k+10)(k-10)$   
**g**  $(2+y)(2-y)$       **h**  $(x+8)(x-8)$   
**i**  $(t+9)(t-9)$
- 2 **a**  $x^2$   
**b i**  $(x-2)$     **ii**  $(x+2)$     **iii**  $x^2$     **iv** 4  
**v**  $x^2-4$   
**c**  $A+B-C=x^2-4$ , which is the area of D, which is  $(x+2)(x-2)$ .
- 3 **a**  $x^2+4x+4-(x^2+2x+1)=2x+3$   
**b**  $(a+b)(a-b)$   
**c**  $(x+2+x+1)(x+2-x-1)=(2x+3)(1)=2x+3$   
**d** The answers are the same.  
**e**  $4x$

### Exercise 9M

- 1  $k = \frac{T}{3}$   
2  $y = X+1$   
3  $p = 3Q$   
4  $r = \frac{A-9}{4}$   
5 **a**  $m = p-t$   
**b**  $t = p-m$
- 6  $m = gv$   
7  $m = \sqrt{t}$   
8  $l = \frac{P-2w}{2}$   
9  $p = \sqrt{m-2}$
- 10 **a**  $-40-32=-72$ ,  $-72 \div 9 = -8$ ,  $5 \times -8 = -40$   
**b**  $68-32=36$ ,  $36 \div 9 = 4$ ,  $4 \times 5 = 20$   
**c**  $f = \frac{9}{5}C + 32$
- 11 Average speeds: outward journey = 72 km/h, return journey = 63 km/h, taking 2 hours. He was held up for 15 minutes.
- 12 **a**  $y = \frac{5x-75}{9}$   
**b** Check by substitution
- 13 **a**  $a = \frac{v-u}{t}$   
**b**  $t = \frac{v-u}{a}$
- 14  $d = \sqrt{\frac{4A}{\pi}}$
- 15 **a**  $y = \frac{x+w}{5}$   
**b**  $w = 5y - x$
- 16  $p = \sqrt{\frac{k}{2}}$
- 17 **a**  $t = u^2 - v$   
**b**  $u = \sqrt{v+t}$
- 18 **a**  $w = k - 5n^2$   
**b**  $n = \sqrt{\frac{K-w}{5}}$
- 19 **a**  $D = \frac{P+Y(K-U)}{3(K-U)}$  or  $\frac{P}{K-U} + \frac{Y}{3}$  or  
 $D = \frac{P}{3(K-U)} + \frac{Y}{3}$   
**b** 16

### Review questions

- 1 **t**  $-xy$
- 2 **a**  $180-p-q$   
**b**  $59^\circ$
- 3 **a**  $r = \frac{C}{2\pi}$   
**b**  $r = \sqrt{\frac{A}{\pi}}$
- 4 **a**  $20x+16$       **b**  $5x+4$
- 5 **a**  $c = \frac{R-3d-5}{7}$       **b** 6
- 6  $12x^2 - xy - 35y^2$
- 7 **a i** 3.5 ml    **ii** 3.7 ml    **iii** 3.84  
**b i** 22    **ii** 38    **iii** 90
- 8  $13.5 \text{ m}^2$
- 9 **a**  $6r(r+h)$     **b**  $h = \frac{A-6r^2}{6r}$     **c** 5 cm
- 10 **b**  $6x^2 - 51x + 90$
- 11 **a**  $x = \frac{A-y}{0.01y}$     **b**  $y(1+0.01x)$     **c** 22 g
- 12 **a i**  $15x^2 - 19x - 56$     **ii**  $16x - 2$   
**b**  $162.25 \text{ cm}^2$
- 13 **a**  $4x^2 + 4x + 1$       **b** 441
- 14 20 m
- 15 **a**  $2(x-8)$     **b**  $x(x-16)$     **c**  $(x-4)(x+4)$   
**d**  $(x-7)(x-9)$
- 16 **a**  $(x-10)(x+10)$   
**b**  $(x-20)(x+20)$   
**c**  $(x-200)(x+200)$
- 17  $-3+7$  is not  $-20$   
 $(x-21)(x+1)$



## Chapter 10 – Ratio and proportion and rates of change: Ratio, speed and proportion

### Exercise 10A

- 1 a 1:3      b 2:3      c 2:5      d 5:8  
e 3:2      f 7:3      g 1:6      h 5:3
- 2 a 1:3      b 5:12      c 17:15      d 4:1  
e 1:24      f 5:2      g 2:1  
h 31:200

3  $\frac{7}{10}$

4  $\frac{2}{5}$

5 a  $\frac{2}{5}$       b  $\frac{3}{5}$

6 2:1

7 a Fruit crush  $\frac{1}{6}$ , lemonade  $\frac{5}{6}$

Fruit Crush	1.25	1	0.2	0.4	0.5
Lemonade	6.25	5	1	2	2.5

b 0.4 litres      c 2.5 litres

8 Sugar  $\frac{5}{22}$ , flour  $\frac{3}{11}$ , margarine  $\frac{2}{11}$ ,  
fruit  $\frac{7}{22}$

9 4

10 1:4

11 a 5:3:2      b 20 years old

12  $13\frac{1}{2}$  litres

### Exercise 10B

- 1 a 160 g, 240 g      b 80 kg, 200 kg  
c 175 min, 125 min      d £20, £30, £50  
e £36, £60, £144      f 50 g, 250 g, 300 g

2 a 175 women      b 30%

3 a 40%      b 300 kg

4 21 horses

5 a 1:400 000      b 1:125 000  
c 1:250 000      d 1:25 000  
e 1:20 000      f 1:40 000

- 6 a Map A 1:250 000, Map B 1:1 000 000  
b 2 km      c 1.2 cm or  $\frac{6}{5}$  cm  
d 4.8 cm

7 a 1:1.6      b 1:3.25      c 1:1.125  
d 1:1.44      e 1:5.4      f 1:1.5  
g 1:4.8      h 1:42      i 1:1.25

8 Diesel: Petrol = 60:90.  $\frac{1}{5}$  of 60 = 12.  $\frac{4}{9}$  of 90 = 40. Total red cars = 52 which is more than  $150 \div 3 = 50$ , so yes.

9  $13 - 9 = 4$ .  $4 \div 5 = 0.8$ .  $2 \times 0.8 = 1.6$ ,  $9 + 1.6 = 10.6$

10 Athos has 3 more parts than Zena.  $24 \div 3 = 8$ , so 1 part is 8. Zena has 8 marbles.

### Exercise 10C

1 a 3:2      b 32      c 80

2 1000 g or 1 kg

3 8100

4 a 14 minutes      b 75 minutes

5 Kevin £2040, John £2720

- 6 a Lemonade 20 litres, ginger 0.5 litres  
b The one in b. In part a there are 50 parts in the ratio 40:9:1, so ginger is  $\frac{1}{50}$  of total amount; in part b there are 13 parts in the ratio 10:2:1, so ginger is  $\frac{1}{13}$  of total amount.  
 $\frac{1}{13} > \frac{1}{50}$

7 296 seats

8 100 students

9 40 ml

- 10 a Will as his multiple of 10 is also a multiple of 9  
b Zeke has rounded off to 1dp and and Yoko has rounded off to 2 dp. They have not used a recurring decimal notation.

11 43 people (20 boys, 8 girls, 15 adults)

12  $\frac{6}{11}$

13  $\frac{1}{2}$

### Exercise 10D

1 18 mph

2 280 miles

3 52.5 mph

- 4 11.50 am
- 5 a 75 mph    b 6.5 h  
c 175 miles    d 240 km  
e 64 km/h    f 325 km  
g 4.3 h (4 h 18 min)
- 6 a 500 m or 0.5 km  
b 12:06 p.m.  
c His actual speed and distance may not be the same as his estimate, so he may not be back at the estimated time.
- 7 a 2.25 h    b 99 miles
- 8 a 1.25 h    b 1 h    15 min
- 9 a Sheffield to London via Midland mainline 74.38 mph. Sheffield to London via East Coast mainline 78.26  
b Doncaster to London 94.12 mph
- 10 a 120 km    b 48 km/h
- 11 a 10 m/s    b 3.3 m/s    c 16.7 m/s
- 12 a 90 km/h    b 43.2 km/h  
c 1.8 km/h
- 13 18 m/s is 64.8 km/h. 40 km at 64.8 km/h is 0.617 hours  $\approx$  37 minutes so train arrives at 8.07 am

14

Time	Cyclist A	Cyclist B
10	16	0
10.15	20	6
10.30	24	12
10.45	28	18
11	32	24
11.15	36	30
11.30	40	36
11.45	44	42
12	48	48
12.15	52	54

Cyclist B catches cyclist A at 12 noon

- 15 Rebecca: 10 minutes at 50 mph covers 8.333 miles, 10 minutes at 70 mph covers 11.666 miles, so total distance is 20 miles in 20 minutes which is 60 mph, so Rebecca is correct.

Nick: 10 miles at 40 mph takes 15 minutes, 10 miles at 60 mph takes 10 minutes, so total distance is 20 miles in 25 minutes, which is 48 mph, so Nick is wrong.

- 16 Josh should take 40 minutes. Nell should take  $50 \div 70 \times 60 = 43$  minutes, but Josh is likely to meet traffic through town so is unlikely to travel at anywhere near 30 mph. Nell is likely to be able to travel at 70 mph on the motorway.

### Exercise 10E

- 1 60 g
- 2 £5.22
- 3 45 trees
- 4 £6.72
- 5 a £312.50    b 8 textbooks
- 6 a 56 litres    b 350 miles
- 7 a i 100 g, 200 g, 250 g, 150 g  
ii 150 g, 300 g, 375 g, 225 g  
iii 250 g, 500 g, 625 g, 375 g
- b 24 biscuits
- 8 a Peter:  $\pounds 2.30 \div 6 = 38.33\text{p}$  each; I can buy four packs (24 sausages) from him (£9.20)  
Paul:  $\pounds 3.50 \div 10 = 35\text{p}$  each; I can only buy two packs (20 sausages) from him (£7)  
I should use Peter's shop to get the most sausages for £10.
- b Buy one pack from Peter and two packs from Paul. That would amount to 26 sausages for £9.30.
- 9  $400 \div 10 = 40$  loaves.  $40 \div 3 = 13.333$ , so  $13.333 \times 1.8 = 24$  kg of flour.

- 10 4 buns and 5 cakes

### Exercise 10F

- 1 a £4.50 for a 10-pack  
b £1.08 for 6  
c £2.45 for 1 litre  
d Same value
- 2 a Large jar as more g per £  
b 75 ml tube as more ml per £  
c Large box as more g per £  
d 400 ml bottle as more ml per £

- 3 a £5.11  
b Large tin (small £5.11/l, medium £4.80/l, large £4.47/l)
- 4 a 95p      b Family size
- 5 Mary
- 6 Kelly
- 7 12-pack  $360 \div 12 = 30$  p per sachet. 40-pack  $1150 \div 40 = 28.75$  p per sachet. 4 sachets cost  $4 \times 35 = \text{£}1.40$  but you get 5, so  $140 \div 5 = 28$  p per sachet, so the offer is the best value.
- 8 a Abe uses  $10 \times 0.75 = 7.5$  litres to do 100 km. Caryl uses  $100 \div 14 = 7.14$  litres to do 100 km and Des uses  $100 \div 55 \times 4.55 = 8.27$  litres to do a 100 km, so Caryl's car is the most economical.  
b It does not give a 'unit' value, i.e. miles per gallon or litres per mile.
- 9 Yes. Let the initial price be  $x$ . 50% off means the cost is  $0.5x$  each. Two for the price of one means two will cost  $x$ , so one will cost  $0.5x$ .
- 10  $\$2.09 \div 1.51 = \text{£}1.38$  per US gallon.  
 $\text{£}1.38 \div 3.8 = \text{£}0.36$  per litre in US.  
 $\text{£}1.09 \div \text{£}0.36 = 3.02$ .  
Petrol is therefore approximately 3 times more expensive in UK than in the USA.
- 11  $\text{£}42 \times 1.2 = \text{£}50.40$   
 $65 \div 50.4 = 1.28$   
 $\text{€}1.28 = \text{£}1$
- 12 a Shirts 4 U:  $75 + 400 \times 3.45 = \text{£}1455$   
Top Tees:  $100 + 200 \times 3.99 + 200 \times 2.75 = \text{£}1448$   
Therefore Top Tees is cheaper.  
b Shirts 4 U:  $75 + 3.45 \times 180 = \text{£}1386$   
Top Tees:  $100 + 200 \times 3.99 + 180 \times 2.75 = \text{£}1393$   
Therefore Shirts 4 U is cheaper.

- 13 a Yogurt  $3 \times 250 = 750$ , Banana  $5 \times 150 = 750$  Kiwi  $17 \times 90 = 1530$   
Ratio  $1530 : 750 : 750 \approx 1500 : 750 : 750 = 2 : 1 : 1$   
This is a sensible recipe because the ratio means they can use whole fruits and whole cartons of yoghurt.  
b 15 Kiwi 1350 g, 2 banana 300 g, 4 yogurt 1000 g  
 $1350 : 300 : 1000 \approx 1333 : 333 : 1000 = 4 : 1 : 3$   
Other answers are possible.

## Review questions

- 1 £3.20
- 2 a 150 g      b 472.5 ml
- 3 48 mph
- 4 225 km
- 5 72 km/h
- 6 Totals are 40 and 60 giving 2 : 3 and a total of 100. 9 : 11 is a ratio of 45 : 55 so swap 10 and 15
- 7 a  $1.73 \div 0.04 = 43.25$  so 43 horses,  $2.64 \div 0.065 = 40.61$  so 40 cattle and  $0.95 \div 0.01 = 95$ . Total  $43 + 40 + 95 = 178$  animals.  
b Horses in field A = 43, Sheep in field B =  $2.64 \div 0.01 = 264$ , Cattle in field C,  $0.95 \div 0.065 = 14.61$ , so 14 cattle.  
Total  $43 + 264 + 14 = 321$  animals.
- 8 3 : 2
- 9 65 blue counters
- 10 a 22.5 kg    b 30 kg    c £19.80
- 11 a  $42 : 63 = 2 : 3$       b 60 out of 200 = 30%  
c

	Men	Women
40 years or over	70	54
Under 40 years	54	72

## Chapter 11 – Geometry and measures: Perimeter and area

### Exercise 11A

- 1 a 24 cm, 35 cm<sup>2</sup>      b 28 cm, 33 cm<sup>2</sup>  
 c 36 cm, 45 cm<sup>2</sup>      d 34 cm, 70 cm<sup>2</sup>  
 e 28 cm, 48 cm<sup>2</sup>      f 14 cm, 10 cm<sup>2</sup>  
 g 26 cm, 42 cm<sup>2</sup>      h 36 cm, 77 cm<sup>2</sup>
- 2 a 20 cm, 24 cm<sup>2</sup>      b 36 cm, 80 cm<sup>2</sup>  
 c 3 cm, 26 cm      d 7 cm, 20 cm<sup>2</sup>  
 e 3 cm, 18 cm<sup>2</sup>      f 6 cm, 30 cm<sup>2</sup>
- 3 a 530 m      b 17400 m<sup>2</sup>
- 4  $160 \times 70 = 11200$ ,  $11200 \div 10000 = 1.12$  hectares
- 5 19 bags

### Exercise 11B

- 1 a 10 cm      b 12 cm      c 12 cm  
 d 14 cm      e 12 cm      f 12 cm
- 2 18 m
- 3 No, the perimeter is 30 cm as two sides will touch.
- 4 False, it is 28 cm. The sum of the lengths of all the sides is 28 cm.
- 5 a 30 cm<sup>2</sup>      b 40 cm<sup>2</sup>      c 51 cm<sup>2</sup>  
 d 35 cm<sup>2</sup>      e 43 cm<sup>2</sup>      f 51 cm<sup>2</sup>  
 g 48 cm<sup>2</sup>      h 33 cm<sup>2</sup>
- 6 24
- 7 The correct answer is 44 cm<sup>2</sup>, the length of the bottom rectangle is 6 cm (10 – 4).
- 8 72 cm<sup>2</sup>
- 9 48 cm
- 10 Yes, the area to paint is 9.1 m<sup>2</sup>

### Exercise 11C

- 1 a 6 cm<sup>2</sup>, 12 cm      b 120 cm<sup>2</sup>, 60 cm  
 c 30 cm<sup>2</sup>, 30 cm
- 2 40 cm<sup>2</sup>
- 3 84 m<sup>2</sup>
- 4 a 21 cm<sup>2</sup>      b 55 cm<sup>2</sup>      c 165 cm<sup>2</sup>
- 5 c 75 cm<sup>2</sup>

- 6 May is correct with 32 cm, but Maggie is incorrect, the area should be 36 cm<sup>2</sup>, it looks like she has forgotten to halve her calculations.
- 7 Area of rectangle =  $5 \times 12 = 60$  cm<sup>2</sup>. The two triangles are the same with one side 6 cm (half of 12) and the other half of (21 – 5) which is 8 cm. Area of one triangle =  $\frac{1}{2} \times 6 \times 8 = 24$  cm<sup>2</sup>. Area of two triangles = 48 cm<sup>2</sup>, so total area =  $60 + 48 = 108$  cm<sup>2</sup>.
- 8 a 5.5 m<sup>2</sup>      b 4

### Exercise 11D

- 1 a 21 cm<sup>2</sup>      b 12 cm<sup>2</sup>      c 14 cm<sup>2</sup>  
 d 55 cm<sup>2</sup>      e 90 cm<sup>2</sup>      f 140 cm<sup>2</sup>
- 2 64 cm<sup>2</sup>
- 3 a 40 cm<sup>2</sup>      b 65 m<sup>2</sup>      c 80 cm<sup>2</sup>
- 4 a 65 cm<sup>2</sup>      b 50 m<sup>2</sup>
- 5 For example: height 10 cm, base 10 cm; height 5 cm, base 20 cm; height 25 cm, base 4 cm; height 50 cm, base 2 cm
- 6 a 1500 cm<sup>2</sup>      b 1800 cm<sup>2</sup>
- 7 Triangle c; a and b each have an area of 15 cm<sup>2</sup> but c has an area of 16 cm<sup>2</sup>.
- 8 For example: Let vertical side be  $T$ , then area =  $\frac{1}{2} \times 2 \times T = T$  cm<sup>2</sup>. The perimeter is  $2 + T + \text{other side}$ , which is bigger than  $T$ . So Tim is correct.

### Exercise 11E

- 1 a 96 cm<sup>2</sup>      b 70 cm<sup>2</sup>      c 20 cm<sup>2</sup>  
 d 125 cm<sup>2</sup>      e 10 cm<sup>2</sup>      f 112 m<sup>2</sup>
- 2 No, it is 24 cm<sup>2</sup>, she used the slanting side instead of the perpendicular height.
- 3 For example: If  $x = 20$ , then the base of each parallelogram is  $\frac{1}{2}(20 - 4) = 8$ . This would make the area of each parallelogram  $8 \times 5 = 40$ , making the whole shape  $4 \times 40 = 160$ , which is incorrect. The correct value of  $x$  is 16.
- 4 a 500 cm<sup>2</sup>      b 15

**Exercise 11F**

- 1 a  $30 \text{ cm}^2$  b  $77 \text{ cm}^2$  c  $24 \text{ cm}^2$   
 d  $42 \text{ cm}^2$  e  $40 \text{ m}^2$  f  $6 \text{ cm}$   
 g  $3 \text{ cm}$  h  $10 \text{ cm}$
- 2 a  $27.5 \text{ cm}, 36.25 \text{ cm}^2$  b  $33.4 \text{ cm}, 61.2 \text{ cm}^2$   
 c  $38.6 \text{ m}, 88.2 \text{ m}^2$
- 3 Any pair of lengths that add up to  $10 \text{ cm}$   
 For example:  $1 \text{ cm}, 9 \text{ cm}; 2 \text{ cm}, 8 \text{ cm}; 3 \text{ cm}, 7 \text{ cm}; 4 \text{ cm}, 6 \text{ cm}; 4.5 \text{ cm}, 5.5 \text{ cm}$
- 4 Shape c. Its area is  $25.5 \text{ cm}^2$
- 5 Shape a. Its area is  $28 \text{ cm}^2$
- 6  $45 \text{ cm}^2$
- 7  $2 \text{ cm}$
- 8  $1.4 \text{ m}^2$
- 9 b  $64 \text{ cm}^2$

**Exercise 11G**

- 1 Top of head-tangent, eyebrows-arcs, nose-sector, mouth-segment, face-circle
- 2 **Diameter** The width of a circle  
**Radius** The distance from the centre to the edge of the circle  
**Circumference** The perimeter of a circle  
**Chord** A line joining two points on a circumference  
**Tangent** A line that touches the circumference of the circle at one point only  
**Arc** A part of the circumference of a circle  
**Sector** A part of a circle lying between two radii and an arc  
**Segment** A part of the area of a circle, lying between a chord and an arc

- 3 Students' own drawings
- 4 Students' own drawings

**Exercise 11H**

- 1 a  $25.1 \text{ cm}$  b  $15.7 \text{ cm}$  c  $44.0 \text{ cm}$   
 d  $22.0 \text{ cm}$  e  $18.8 \text{ cm}$  f  $47.1 \text{ cm}$   
 g  $28.9 \text{ cm}$  h  $14.8 \text{ cm}$

- 2 a  $6.3 \text{ cm}$  b  $8.2 \text{ cm}$  c  $5.3 \text{ cm}$   
 d  $7.5 \text{ cm}$
- 3 a  $31.4 \text{ cm}$  b  $18.8 \text{ cm}$  c  $9.4 \text{ cm}$   
 d  $25.1 \text{ cm}$  e  $5.7 \text{ cm}$  f  $15.7 \text{ cm}$   
 g  $81.7 \text{ cm}$  h  $39.6 \text{ cm}$
- 4 a  $198 \text{ cm}$  b  $505$
- 5 a A  $188.5 \text{ m}$ , B  $194.8 \text{ m}$ , C  $201.1 \text{ m}$ , D  $207.3 \text{ m}$   
 b  $18.7$  or  $18.8 \text{ m}$
- 6  $879.6$  or  $880 \text{ cm}$
- 7 a  $37.7 \text{ cm}$  b  $3770 \text{ cm}$  c  $37.7 \text{ m}$   
 d  $37.7 \text{ km}$
- 8  $100 \text{ cm}$
- 9  $24.2 \text{ cm}$
- 10  $15.9 \text{ cm}$
- 11  $25.7 \text{ cm}$
- 12 a Fir:  $62.8 \text{ cm}$ , ash:  $75.4 \text{ cm}$ , elm:  $69.1 \text{ cm}$ , oak:  $81.7 \text{ cm}$   
 b The difference between the distances round the circumferences of two trees is  $2\pi$  times the difference between their radii, or  $\pi$  times the difference between their diameters.  
 c  $6.28 \text{ m}$
- 13 a Perimeters of shapes A and B are both  $25.1 \text{ cm}$   
 b  $25.1 \text{ cm}$
- 14  $4a = 2\pi r$ , so  $2a = \pi r$ , therefore  $r = \frac{2a}{\pi}$
- 15  $7$
- 16 a  $4\pi = 12.6 \text{ cm}$  b New radius of small circle will be  $3$ , giving new perimeter of the whole shape of  $12\pi$ , which is three times larger than  $4\pi$ .

**Exercise 11I**

- 1 a  $78.5 \text{ cm}^2$  b  $28.3 \text{ cm}^2$  c  $7.1 \text{ cm}^2$   
 d  $50.3 \text{ cm}^2$  e  $2.5 \text{ cm}^2$  f  $19.6 \text{ cm}^2$   
 g  $530.9 \text{ cm}^2$  h  $124.7 \text{ cm}^2$
- 2 a  $3.1 \text{ cm}^2$  b  $5.3 \text{ cm}^2$  c  $2.3 \text{ cm}^2$   
 d  $4.5 \text{ cm}^2$
- 3 a  $50.3 \text{ cm}^2$  b  $19.6 \text{ cm}^2$  c  $153.9 \text{ cm}^2$   
 d  $38.5 \text{ cm}^2$  e  $28.3 \text{ cm}^2$  f  $176.7 \text{ cm}^2$   
 g  $66.5 \text{ cm}^2$  h  $17.3 \text{ cm}^2$

- 4 a  $9.1 \text{ cm}^2$  c 138 b  $2000 \text{ cm}^2$   
 d  $1255.8 \text{ cm}^2$  or  $1252.9 \text{ cm}^2$  using unrounded answer from a  
 e  $744.2 \text{ cm}^2$  or  $747.1 \text{ cm}^2$ , using unrounded answer from a
- 5  $3848.5 \text{ m}^2$
- 6 a i  $56.5 \text{ cm}$  ii  $254.5 \text{ cm}^2$   
 b i  $69.1 \text{ cm}$  ii  $380.1 \text{ cm}^2$   
 c i  $40.8 \text{ cm}$  ii  $132.7 \text{ cm}^2$   
 d i  $88.0 \text{ cm}$  ii  $615.8 \text{ cm}^2$
- 7  $962.9 \text{ cm}^2$  (or  $962.1 \text{ cm}^2$ )
- 8 a  $56.5 \text{ cm}^2$  b  $19.6 \text{ cm}^2$
- 9 a  $50.3 \text{ m}^2$  b  $44.0 \text{ cm}^2$
- 10  $a^2 = \pi r^2$ , so  $r^2 = \frac{a^2}{\pi}$ , therefore  $r = \frac{a}{\sqrt{\pi}}$
- 11 Area of the square is  $100 \text{ cm}^2$ , Area of circle is  $\pi \times 5^2 = 78.54$ , leaving the shaded area as  $100 - 78.54 = 21.46 \text{ cm}^2$ . Percentage of shaded area =  $\frac{21.46}{100} \times 100 = 21.46\%$ , which rounds to  $21\%$ , so Lewis is correct.
- 12 a  $4\pi$  or  $12.6 \text{ cm}^2$   
 b i  $16\pi$  or  $50.3 \text{ cm}^2$  ii  $36\pi$  or  $113 \text{ cm}^2$   
 iii  $64\pi$  or  $201 \text{ cm}^2$   
 c  $a = 4\pi r^2$

### Exercise 11J

- 1 a  $4\pi \text{ cm}$  b  $20\pi \text{ cm}$  c  $15\pi \text{ cm}$   
 d  $4\pi \text{ cm}$
- 2 a  $16\pi \text{ cm}^2$  b  $25\pi \text{ cm}^2$  c  $9\pi \text{ cm}^2$   
 d  $81\pi \text{ cm}^2$
- 3  $25 \text{ cm}$
- 4  $10 \text{ cm}$
- 5 Rory is correct. Area =  $\pi \times \left(\frac{100}{\pi}\right)^2$ , so Area =  $\pi \times \frac{100^2}{\pi^2} = \frac{10000}{\pi}$ , so Alice was incorrect. She needed to put an extra zero in the numerator.
- 6 a  $12.5\pi \text{ cm}^2$  b  $16\pi \text{ cm}^2$  c  $(16\pi + 80) \text{ cm}^2$   
 d  $(50\pi + 100) \text{ cm}^2$
- 7 a  $32\pi \text{ cm}^2$  b  $16\pi \text{ cm}^2$   
 c  $8\pi \text{ cm}^2$  d  $4\pi \text{ cm}^2$
- 8  $18\pi - 9\pi = 9\pi$
- 9 c;  $80 + \frac{1}{4}(\pi \times 8^2) = 80 + 16\pi$


## Review questions

- 1  $30 \text{ cm}^2$
- 2 48 tiles
- 3 Area =  $\frac{1}{2} \times 12 \times 9 = 54 \text{ cm}^2$ , Tilly has forgotten the  $\frac{1}{2}$  in her calculation.
- 4  $8 \text{ cm}$
- 5 209 sheep
- 6  $\pi d = 30$ , so  $d = \frac{30}{\pi}$ , so radius =  $\frac{15}{\pi}$ , so area =  $\pi \times \left(\frac{15}{\pi}\right)^2 = \frac{225}{\pi} = 71.6$
- 7 Circle area =  $\pi \times 25 = 78.5$ , Parallelogram area =  $9 \times 8 = 72$ , difference =  $78.5 - 72 = 6.5$ , Andy is correct.
- 8 a  $2.5 \text{ cm}$   
 b Area of large circle =  $10^2 \times \pi = 100\pi$ , area of two small circles is  $2 \times \pi \times 2.5^2 = 12.5\pi$ , the shaded part =  $(100 - 12.5)\pi = 87.5\pi$ .  
 The percentage of shaded =  $\frac{87.5\pi}{100\pi} \times 100 = 87.5\%$
- 9  $\frac{20}{\pi}$
- 10  $25.7 \text{ cm}$

## Chapter 12 – Geometry and measures: Transformations

### Exercise 12A

- 1 a 4 b 5 c 6  
 d 4 e 6
- 2 a 2 b 2 c 2  
 d 2 e 2
- 3 A, B, C, D, E, F, G, J, K, L, M, P, Q, R, T, U, V, W, Y
- 4 for example:



5

		Number of lines of symmetry			
		0	1	2	3
Order of rotational symmetry	1	D	A		
	2	E		B	
	3				C

6 She is correct since the angle sum around the centre point is 360 and  $360 \div 3 = 120$

7 Yes she is correct. A triangle can only have 1 or 3 lines of symmetry. If a triangle has 3 lines of symmetry it also has rotational symmetry of order 3, so this triangle must only have 1 line of symmetry. This will mean it has two angles identical and two equal sides, and hence an isosceles triangle.

### Exercise 12B

1 Check students' translations.

2 Check students' translations.


3 a i  $\begin{pmatrix} 1 \\ 3 \end{pmatrix}$  ii  $\begin{pmatrix} 4 \\ 2 \end{pmatrix}$  iii  $\begin{pmatrix} 2 \\ -1 \end{pmatrix}$  iv  $\begin{pmatrix} 5 \\ 1 \end{pmatrix}$

b i  $\begin{pmatrix} -1 \\ -3 \end{pmatrix}$  ii  $\begin{pmatrix} 3 \\ -1 \end{pmatrix}$  iii  $\begin{pmatrix} -2 \\ 3 \end{pmatrix}$  iv  $\begin{pmatrix} 3 \\ 3 \end{pmatrix}$

c i  $\begin{pmatrix} -2 \\ -3 \end{pmatrix}$  ii  $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$  iii  $\begin{pmatrix} -5 \\ 4 \end{pmatrix}$  iv  $\begin{pmatrix} 0 \\ 4 \end{pmatrix}$

d i  $\begin{pmatrix} 3 \\ 2 \end{pmatrix}$  ii  $\begin{pmatrix} -4 \\ 2 \end{pmatrix}$  iii  $\begin{pmatrix} 5 \\ -4 \end{pmatrix}$  iv  $\begin{pmatrix} -2 \\ -7 \end{pmatrix}$

4



5 a  $\begin{pmatrix} -3 \\ -1 \end{pmatrix}$  b  $\begin{pmatrix} 4 \\ -4 \end{pmatrix}$  c  $\begin{pmatrix} -5 \\ -2 \end{pmatrix}$

d  $\begin{pmatrix} 4 \\ 7 \end{pmatrix}$  e  $\begin{pmatrix} -1 \\ 5 \end{pmatrix}$  f  $\begin{pmatrix} 1 \\ 6 \end{pmatrix}$

g  $\begin{pmatrix} -4 \\ 4 \end{pmatrix}$  h  $\begin{pmatrix} -4 \\ -7 \end{pmatrix}$

6  $10 \times 10 = 100$  (including  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ )

7 a Check students' designs for a Snakes and ladders board.

b because the ladders always mean moving up the board

ii the snakes always mean moving down the board

8  $\begin{pmatrix} -x \\ -y \end{pmatrix}$


9 a Check student's diagram b  $\begin{pmatrix} -300 \\ -500 \end{pmatrix}$

10 a Check student's diagram b  $\begin{pmatrix} -1 \\ 4 \end{pmatrix}$


11 Multiply each part of the vector describing the move by  $-1$

### Exercise 12C


1 a














b




c






**4 a-b**




**5 a-e**




**f** Reflection in the y-axis

**6 a-b**



**c** y-value changes sign  
**d**  $(a, -b)$

**7 a-b**




**c** x-value changes sign  
**d**  $(-a, b)$




8 Possible answer: Take the centre square as ABCD then reflect this square each time in the line, AB, then BC, then CD and finally AD.

9  $x = -1$




11 a-i




j A reflection in  $y = x$

12 a-c




13 a-c



d Coordinates are reversed:  $x$  becomes  $y$  and  $y$  becomes  $x$

e  $(b, a)$

14 a-c




d Coordinates are reversed and change sign,  $x$  becomes  $-y$  and  $y$  becomes  $-x$

e  $(-b, -a)$


15 A reflection in the same line

### Exercise 12D


1 a




b




c




d





2 a





b






**3 a**




- b i** Rotation  $90^\circ$  anticlockwise  
**ii** Rotation  $90^\circ$  clockwise

**4 a-d**




**5**



- b** Rotate shape  $60^\circ$  around point B, then repeat another four times.

- 6** Possible answer: If ABCD is the centre square, rotate about A  $90^\circ$  anticlockwise, then rotate about A  $180^\circ$ , now rotate about new B  $180^\circ$ , and finally rotate about new C  $180^\circ$ .

**7**




- 8 a** Because the opposite rotations equal a total of  $360^\circ$ , which is the same point

- b i**  $90^\circ$  anticlockwise  
**ii**  $270^\circ$  anticlockwise  
**iii**  $300^\circ$  clockwise  
**iv**  $260^\circ$  clockwise

- 9** Show by drawing a shape or use the fact that  $(a, b)$  becomes  $(a, -b)$  after reflection in the  $x$ -axis, and  $(a, -b)$  becomes  $(-a, -b)$  after reflection in the  $y$ -axis, which is equivalent to a single rotation of  $180^\circ$ .

- 10** She is correct.

**11 a**











- b** i Rotation  $60^\circ$  clockwise about O  
 ii Rotation  $120^\circ$  clockwise about O  
 iii Rotation  $180^\circ$  about O  
 iv Rotation  $240^\circ$  clockwise about O  
**c** i Rotation  $60^\circ$  clockwise about O  
 ii Rotation  $180^\circ$  about O

12 Rotation  $90^\circ$  anticlockwise about (3, 22)

13 A rotation of the same angle but in the opposite direction. In this case the direction would be in the clockwise direction.

### Exercise 12E



b 3 : 1

c 3 : 1

d 9 : 1

5 (1, -5), (3, -3), (-5, -5)

6 b Rotation  $180^\circ$  around (-1, 2)

7 a Enlargement by scale factor -2 from centre of enlargement (1, 3)

b Enlargement by scale factor  $-\frac{1}{2}$  from centre of enlargement (1, 3)

### Exercise 12F

1 a (-4, -3) and (-2, -1)


2 a (-5, 2)      b (-5, -2)

c Rotation of  $180^\circ$  about the origin.

3 A: translation  $\begin{pmatrix} 1 \\ -2 \end{pmatrix}$ , B: reflection in y-axis,

C: rotation  $90^\circ$  clockwise about (0, 0), D: reflection in  $x = 3$ , E: reflection in  $y = 4$ , F: enlargement by scale factor 2, centre (0, 1)

4



- 5 a  $T_1$  to  $T_2$ : rotation  $90^\circ$  clockwise about  $(0, 0)$   
 b  $T_1$  to  $T_6$ : rotation  $90^\circ$  anticlockwise about  $(0, 0)$


c  $T_2$  to  $T_3$ : translation  $\begin{pmatrix} 2 \\ 2 \end{pmatrix}$

d  $T_6$  to  $T_2$ : rotation  $180^\circ$  about  $(0, 0)$

e  $T_6$  to  $T_5$ : reflection in  $y$ -axis

f  $T_5$  to  $T_4$ : translation  $\begin{pmatrix} 4 \\ 0 \end{pmatrix}$

6 a-d




e  $T_d$  to  $T$ : rotation  $90^\circ$  anticlockwise about  $(0, 0)$

7  $(3, 1)$

8 For example: Reflection in  $x$ -axis, translation  $\begin{pmatrix} 0 \\ -5 \end{pmatrix}$ , rotation  $90^\circ$  clockwise about  $(0, 0)$

9 For example: Translation  $\begin{pmatrix} 0 \\ -8 \end{pmatrix}$ , reflection in  $x$ -axis, rotation  $90^\circ$  clockwise about  $(0, 0)$

10 a




b enlargement scale factor 2 from  $(0, -2)$


11 No, the two reflections will result in the original orientation.

### Exercise 12G

1



2 a



b i  $\begin{pmatrix} 3 \\ 6 \end{pmatrix}$     ii  $\begin{pmatrix} -2 \\ -9 \end{pmatrix}$     iii  $\begin{pmatrix} 1 \\ -3 \end{pmatrix}$

3 a  $(0, 0)$  and  $(3, 5)$     b  $\begin{pmatrix} 3 \\ 5 \end{pmatrix}$

4 a  $(2, 2), (6, 1), (6, 4)$   
 b  $(-6, -1), (-2, -2), (-2, 1)$

5  $(1, 5)$

6 a  $(3, 8)$     b  $(6, -6)$

## Exercise 12H

- 1 a Any three of:  $\overline{AC}$ ,  $\overline{CF}$ ,  $\overline{BD}$ ,  $\overline{DG}$ ,  $\overline{GI}$ ,  $\overline{EH}$ ,  $\overline{HJ}$ ,  $\overline{JK}$   
 b Any three of:  $\overline{BE}$ ,  $\overline{AD}$ ,  $\overline{DH}$ ,  $\overline{CG}$ ,  $\overline{GJ}$ ,  $\overline{FI}$ ,  $\overline{IK}$   
 c Any three of:  $\overline{AO}$ ,  $\overline{CA}$ ,  $\overline{FC}$ ,  $\overline{IG}$ ,  $\overline{GD}$ ,  $\overline{DB}$ ,  $\overline{KJ}$ ,  $\overline{JH}$ ,  $\overline{HE}$   
 d Any three of:  $\overline{BO}$ ,  $\overline{EB}$ ,  $\overline{HD}$ ,  $\overline{DA}$ ,  $\overline{JG}$ ,  $\overline{GC}$ ,  $\overline{KI}$ ,  $\overline{IF}$


- 2 a  $2a$       b  $2b$       c  $a + b$   
 d  $2a + b$     e  $2a + 2b$     f  $a + 2b$   
 g  $a + b$       h  $2a + 2b$     i  $3a + b$   
 j  $2a$         k  $b$             l  $2a + b$

- 3 a Equal      b  $\overline{AI}$ ,  $\overline{BJ}$ ,  $\overline{DK}$

- 4 a  $\overline{OJ} = 2\overline{OD}$  and parallel    b  $\overline{AK}$

- c  $\overline{OF}$ ,  $\overline{BI}$ ,  $\overline{EK}$

5



- 6 a Lie on same straight line  
 b All multiples of  $a + b$  and start at O  
 c  $4a + 2b$

- 7 a  $-b$       b  $3a - b$       c  $2a - b$   
 d  $a - b$     e  $a + b$         f  $-a - b$   
 g  $2a - b$     h  $-a - 2b$     i  $a + 2b$   
 j  $-a + b$     k  $2a - 2b$     l  $a - 2b$

- 8 a Equal but in opposite directions

- b Any three of:  $\overline{DA}$ ,  $\overline{EF}$ ,  $\overline{GJ}$ ,  $\overline{FI}$ ,  $\overline{AH}$

- 9 a Opposite direction and  $\overline{AB} = -\frac{1}{2}\overline{CK}$

- b  $\overline{BJ}$ ,  $\overline{CK}$


- c  $\overline{EB}$ ,  $\overline{GO}$ ,  $\overline{KH}$

- 10 a i  $3a + 2b$     ii  $3a + b$   
 iii  $2a - b$     iv  $2b - 2a$


- b  $\overline{DG}$  and  $\overline{BC}$

## Review questions

- 1 a 3  
 b There will be 3 lines all through a vertex and the opposite side's midpoint.



- 2 a




- b  $90^\circ$  anticlockwise rotation about  $(0, 0)$

- 3  $180^\circ$  rotation about  $(0, 0)$


- 4 Translation with vector  $\begin{pmatrix} 4 \\ 1 \end{pmatrix}$

- 5 a



- c Rotation of  $180^\circ$  about  $(-1, 1)$

6 a – b



c Rotation of half a turn around the origin

7 (9, 0), (11, -3), (2, -1)

8 i –b ii –a iii b iv a v a + b vi a – b

9 Students' diagram.

10 For example: Reflection in the y axis, reflection in the line  $y = 1$

## Chapter 13 – Probability: Probability and events

### Exercise 13A

1 a unlikely b certain c likely d very unlikely e impossible f very likely g evens

2 a  $\frac{1}{6}$  b  $\frac{1}{6}$  c  $\frac{1}{2}$  d  $\frac{1}{13}$  e  $\frac{1}{4}$   
f  $\frac{1}{2}$  g 1 h  $\frac{1}{26}$  i  $\frac{1}{13}$  j 0

3 a 0 b 1

4 a  $\frac{1}{10}$  b  $\frac{1}{2}$  c  $\frac{2}{5}$  d  $\frac{1}{5}$  e  $\frac{3}{5}$

5 a  $\frac{1}{2}$  b  $\frac{1}{3}$  c  $\frac{1}{6}$

6  $\frac{1}{50}$

7 a  $\frac{1}{13}$  b  $\frac{1}{13}$  c  $\frac{1}{2}$  d  $\frac{1}{26}$

8 a i  $\frac{12}{25}$  ii  $\frac{7}{25}$  iii  $\frac{6}{25}$

b They add up to 1.

c All possible outcomes are used.

9 a AB, AC, AD, AE, BC, BD, BE, CD, CE, DE

b 1 c  $\frac{1}{10}$  d 6 e  $\frac{3}{5}$  f  $\frac{3}{10}$

10 There might not be the same number of boys as girls in the class.

11 Class U

### Exercise 13B

1 a  $\frac{19}{20}$  b 55% c 0.2

2 a i  $\frac{3}{13}$  ii  $\frac{10}{13}$

b i  $\frac{1}{4}$  ii  $\frac{3}{4}$

c i  $\frac{2}{13}$  ii  $\frac{11}{13}$

3 a  $\frac{1}{2}$  b 1 c  $\frac{1}{3}$

4 a  $\frac{6}{11}$  b  $\frac{5}{11}$  c  $\frac{6}{11}$

5 a  $\frac{1}{2}$  b  $\frac{1}{5}$  c  $\frac{4}{5}$  d  $\frac{7}{10}$

6 a  $\frac{7}{15}$  b  $\frac{2}{15}$  c  $\frac{13}{15}$  d  $\frac{8}{15}$

e  $\frac{7}{15}$  f 0

7  $P(\text{Noah will lose}) = \frac{1}{2}$  and  $P(\text{Phoebe will lose}) = \frac{1}{3}$ , so  $P(\text{Noah will not lose}) = \frac{1}{2}$  and  $P(\text{Phoebe will not lose}) = \frac{2}{3}$ . Phoebe has a better chance of not losing as  $\frac{2}{3} > \frac{1}{2}$

8 Because it might be possible for the game to end in a draw.

9 a i  $\frac{1}{4}$  ii  $\frac{3}{4}$

b i  $\frac{3}{11}$  ii  $\frac{8}{11}$

10 35%

11 0.5

### Exercise 13C

1 a  $\frac{1}{6}$  b  $\frac{1}{6}$  c  $\frac{1}{3}$

2 a  $\frac{1}{4}$  b  $\frac{1}{4}$  c  $\frac{1}{2}$

3 a  $\frac{2}{11}$  b  $\frac{4}{11}$  c  $\frac{6}{11}$

4 a  $\frac{1}{3}$  b  $\frac{2}{5}$  c  $\frac{11}{15}$  d  $\frac{11}{15}$  e  $\frac{1}{3}$

- 5 a 0.8      b 0.2
- 6 a  $\frac{17}{20}$       b  $\frac{2}{5}$       c  $\frac{3}{4}$
- 7 Because these are not mutually exclusive events and probability cannot exceed 1.
- 8  $\frac{3}{4}$
- 9 The probability for each day stays the same, at  $\frac{1}{4}$ .
- 10 a The choices of drink and snack are not mutually exclusive.  
 b (C, D), (T, G), (T, R), (T, D), (H, G), (H, R), (H, D)  
 c i (C, D) = 0.12, (T, G) = 0.15, (T, R) = 0.05, (T, D) = 0.3, (H, G) = 0.09, (H, R) = 0.03, (H, D) = 0.18  
 ii The total is 1 as all the outcomes are exhaustive.

**Exercise 13D**

- 1 a B      b B      c C      d A  
 e B      f a      g B      h B
- 2 a 0.2, 0.08, 0.1, 0.105, 0.148, 0.163, 0.1645  
 b 6      c 1      d  $\frac{1}{6}$
- 3 a 0.095, 0.135, 0.16, 0.265, 0.345  
 b 40  
 c No; all numbers should be close to 40.
- 4 a 0.2, 0.25, 0.38, 0.42, 0.385, 0.3974  
 b 79
- 5 a Caryl, threw the greatest number of times.  
 b 0.39, 0.31, 0.17, 0.14  
 c Yes; all the relative frequencies should be close to 0.25.
- 6 Thursday; the day with the highest relative frequency
- 7 The missing top numbers are 4 and 5, the bottom two numbers are both likely to be close to 20.
- 8 Although he might expect the probability to be close to  $\frac{1}{2}$ , giving 500 heads, the actual number of heads is unlikely to be exactly 500, but should be close to it.
- 9 Roza is correct, as the expected numbers are: 50, 12.5, 25, 12.5. Sam has not taken into account the fact that there are four red sectors.

**Exercise 13E**


- 1 a  $\frac{1}{6}$       b 25
- 2 a  $\frac{1}{2}$       b 1000
- 3 a i  $\frac{1}{2}$       ii  $\frac{1}{13}$       iii  $\frac{1}{4}$       iv  $\frac{1}{52}$   
 b i 260      ii 40      iii 130      iv 10
- 4 a  $\frac{1}{37}$       b 5
- 5 a 150      b 100      c 250      d 0
- 6 a 167      b 833
- 7 a 0.6      b 120
- 8 a

Score	1	2	3	4	5	6
Expected occurrences	10	10	10	10	10	10

- b
- $$\frac{1 \times 10 + 2 \times 10 + 3 \times 10 + 4 \times 10 + 5 \times 10 + 6 \times 10}{60} = \frac{210}{60} = 3.5$$
- c Find the average of the scores, which is  $\frac{21}{6} = 3.5$
- 9 a 0.111      b 40
- 10 281 days
- 11 Multiply the number of plants by 0.997.
- 12 400
- Exercise 13F**
- 1 a 48      b 12
- 2 a 40
- 3 a (BS, BT), (BS, RT), (BS, GT), (WS, BT), (WS, RT), (WS, GT), (GS, BT), (GS, RT), (GS, GT)  
 b  $\frac{7}{9}$
- 4  $10^9 = 1\,000\,000\,000$  (1 billion)
- 5 a  $\frac{1}{2}$   
 b HHH, HHT, HTH, HTT, THH, THT, TTH, TTT  $\frac{3}{8}$
- 6 20
- 7  $2^5 = 32$
- 8 56

- 9  $26^2 \times 10^2 = 67\,600 > 50\,000$
- 10 a  $2 \times 26 \times 999\,999 \times 4 = 207\,999\,792$   
 b  $\frac{1}{4}$
- 11 a 24                                      b  $\frac{1}{6}$
- 12  $\frac{11}{21}$  Number of possible permutations is 21, £2 + any coins = 6, £1 + any coin other than £2 = 5, etc, so total is 6 + 5 + 4 + 3 + 2 + 1 = 21, Only the ones with a £2 or a £1 in will give a total greater than £1 = 11 ways
- 13 a 6, 16, etc.. up to 196 which is 20 plus 60, 41 up to 69, which is 9 (66 already counted) plus 160, 162 up to 169 which is 9 (166 already counted) giving a total of 38  
 b  $\frac{38}{200} = \frac{19}{100}$   
 c  $\frac{11}{200}$
- 14 a Key pad A,  $10^4 = 10000$  Keypad B  $13^4 = 28561$   
 b  $3 \times 10^3 = 3\,000$                       c  $\frac{1}{24}$
- 15 a i 5040                                      ii  $2.43 \times 10^{18}$  (3 sf)  
 b This depends on your calculator but  $69! = 1.71 \times 10^{98}$ , which is about the number of atoms in 1 quintillion universes. (Look this term up if you're not sure what it means).

### Review questions

- 1 a April and May  
 b Daffodil  
 c February  
 d Crocus  
 e i  $\frac{1}{5}$   
 ii 
- 2  $\frac{5}{12}$
- 3 a  $\frac{13}{50}$                       b  $\frac{1}{5}$                       c  $\frac{1}{2}$                       d  $\frac{7}{10}$
- 4 a 1  
 b There are five possible outcomes and only two of them are 2, so the probability is  $\frac{2}{5}$ .
- 5 a 0.5  
 b yes, as you would expect all the probabilities to be close to 0.25

- 6 a  $\frac{3}{8}$     b 12
- 7 a i 0.8    ii 12  
 b No, as  $P(\text{six}) = \frac{1}{6}$ , so Tom is likely to throw 10 sixes

8 a

Colour	White (W)	Green (G)	Blue (B)
Relative frequency	$\frac{1}{4}$	$\frac{9}{20}$	$\frac{3}{10}$

- b The second table, as there were more trials
- 9 a (H, 1), (H, 2), (H, 3), (H, 4), (H, 5), (H, 6), (T, 1), (T, 2), (T, 3), (T, 4), (T, 5), (T, 6)  
 b  $\frac{2}{12} = \frac{1}{6}$  as (H, 3) scores 6 and (T, 6) scores 6
- 10  $\frac{3}{6} = \frac{1}{2}$ . There are 6 combinations, 2 + 14 = 16, 11 + 14 = 25 and 14 + 22 = 36 give square numbers
- 11 a  $10^3 = 1000$                                       b  $\frac{1}{100}$   
 c Trial and improvement  $9^5 = 59\,049$

## Chapter 14 – Geometry and measures: Volumes and surface areas of prisms

### Exercise 14A

1 a

Name	Number of faces (F)	Number of vertices (V)	Number of edges (E)
Cuboid	6	8	12
Square-based pyramid	5	5	8
Tetrahedron	4	4	6
Octahedron	8	6	12
Triangular prism	5	6	9
Hexagonal prism	8	12	18
Hexagon based pyramid	7	7	12

b  $F + V = E + 2$



- 2 a i 6, 12      ii 24, 96  
 b i 2            ii 4            iii 8  
 c ii 2, 4        ii 18, 108  
 d i 3            ii 9            iii 27
- 3 She might have calculated this by multiplying 5 by 5 by 4 to get 100.
- 4  $65 \text{ cm}^2$
- 5 6
- 6 David is correct. A pentagon has 5 sides, so a pentagonal prism has  $5 + 5 + 5$  edges, and  $2 + 5$  faces.  
 $5 + 5 + 5 + 2 + 5 = 22$ .


**Exercise 14B**

- 1 a i  $198 \text{ cm}^3$             ii  $234 \text{ cm}^2$   
 b i  $90 \text{ cm}^3$             ii  $146 \text{ cm}^2$   
 c i  $1440 \text{ cm}^3$             ii  $792 \text{ cm}^2$   
 d i  $525 \text{ cm}^3$             ii  $470 \text{ cm}^2$
- 2 24 litres
- 3 a  $160 \text{ cm}^3$     b  $480 \text{ cm}^3$     c  $150 \text{ cm}^3$
- 4 a i  $64 \text{ cm}^3$             ii  $96 \text{ cm}^2$   
 b i  $343 \text{ cm}^3$             ii  $294 \text{ cm}^2$   
 c i  $1000 \text{ mm}^3$             ii  $600 \text{ mm}^2$   
 d i  $125 \text{ m}^3$             ii  $150 \text{ m}^2$   
 e i  $1728 \text{ m}^3$             ii  $864 \text{ m}^2$
- 5 86
- 6 a  $180 \text{ cm}^3$     b 5 cm      c 6 cm  
 d 10 cm      e  $81 \text{ cm}^3$
- 7  $32\,000$  litre is equivalent to  $32\,000 \times 1000 \text{ cm}^3 = 32\,000\,000 \text{ cm}^3$ .  $V = lwh$ , so  $650 \times 310 \times h = 32\,000\,000$ , so  $h = \frac{32\,000\,000}{(650 \times 310)} = 158.8 \text{ cm}$ .
- 8 Volume = base area  $\times$  height, so base area = volume/height =  $\frac{168}{3.5} = 48 \text{ m}^2$ . So Peter is only correct if he has rounded to the nearest ten.
- 9 a 3 cm    b 5 m    c 2 mm    d 1.2 m
- 10 a  $148 \text{ cm}^3$             b  $468 \text{ cm}^3$
- 11 If this was a cube, the side length would be 5 cm, so total surface area would be  $5 \times 5 \times 6 = 150 \text{ cm}^2$ ; so this particular cuboid is not a cube.
- 12  $600 \text{ cm}^2$

**Exercise 14C**

- 1 a i  $21 \text{ cm}^2$             ii  $63 \text{ cm}^3$   
 b i  $48 \text{ cm}^2$             ii  $432 \text{ cm}^3$   
 c i  $36 \text{ m}^2$             ii  $324 \text{ m}^3$

- 2 a  $432 \text{ m}^3$     b  $225 \text{ m}^3$     c  $1332 \text{ m}^3$



- b About  $3\frac{1}{2}$  hours
- 4  $V = (1.5 + 3) \div 2 \times 1.7 \times 2 = 7.65 \text{ m}^3$
- 5  $27 = 3 \times 3 \times 3$ ,  $27 + 37 = 64 = 4^3$ ,  $4 - 3 = 1$ . Hence the side length is 1 small cube longer, hence 2 cm longer
- 6 a i  $21 \text{ cm}^3$             ii  $54 \text{ cm}^2$   
 b i  $210 \text{ cm}^3$             ii  $270 \text{ cm}^2$
- 7 10.2 tonnes
- 8 She was silly because 26 880 is simply all the dimensions multiplied together. The volume is  $672 \text{ cm}^3$ .

**Exercise 14D**

- 1 a i  $226 \text{ cm}^3$             ii  $207 \text{ cm}^2$   
 b i  $14.9 \text{ cm}^3$             ii  $61.3 \text{ cm}^2$   
 c i  $346 \text{ cm}^3$             ii  $275 \text{ cm}^2$   
 d i  $1060 \text{ cm}^3$             ii  $636 \text{ cm}^2$
- 2 a  $251 \text{ cm}^3$             b  $445 \text{ cm}^3$   
 c  $2150 \text{ cm}^3$             d  $0.25 \text{ m}^3$
- 3 Volume =  $\pi \times (0.3)^2 \times 4.2 = 0.378\pi$   
 Cost =  $0.378 \times \pi \times \text{£}67.50 = \text{£}80.16$  which is  $\text{£}80$  to 2 sf
- 4 1.23 tonnes
- 5  $665 \text{ cm}^3$
- 6 Label should be less than 10.5 cm wide so that it fits the can and does not overlap the rim and more than 23.3 cm long to allow an overlap.
- 7 Volume =  $\pi \times 32.5^2 \times 1000 = 331\,831 \text{ cm}^3$   
 $1 \text{ litre} = 1000 \text{ cm}^3$   
 volume =  $331\,831 \div 1000 = 331.831 \text{ litres} = 332 \text{ litres}$  (3 sf)
- 8 There is no right answer. Students could start with the dimensions of a real can. Often drinks cans are not exactly cylindrical. One possible answer is height of 6.6 cm and diameter of 8 cm.
- 9 About 127 cm
- 10 Volume of A =  $2^2 \times \pi \times 5 = 20\pi$ , volume of B =  $1^2 \times \pi \times 21 = 21\pi$ , B has the larger volume.
- 11  $146 \text{ cm}^3$

12  $78 \text{ m}^3$  ( $78.3 \text{ m}^3$ )

13 327 litres

## Review questions

1 a  $60 \text{ cm}^3$

b  $104 \text{ cm}^2$

2  $38 \text{ cm}^2$

3 Arrange small boxes so that the 8 by 8 part fills one layer in the 32 by 32 box, giving  $4 \times 4$  boxes. Then this will give 5 layers of  $4 \times 4$  boxes giving  $5 \times 4 \times 4 = 80$  boxes.

4 Area of circular end  $= \pi r^2 = \pi \times 4.5^2 = 63.617251$ , curved surface area  $= \pi dh = \pi \times 9 \times 21 = 593.76101$ , total area  $= 2 \times 63.617251 + 593.76101 = 720.99551 = 721 \text{ cm}^2$  (nearest whole number)

5  $5740 \text{ cm}^3$  (to 3 sf)

6 610 g (2 sf)

7  $600 \text{ cm}^3$

8 360 g

## Chapter 15 – Algebra:

### Linear equations

#### Exercise 15A

1 a  $x = 4$

b  $w = 14$

c  $x = 5$

d  $x = 6$

e  $z = 24$

f  $x = 2.5$

2 Any valid equation such as  $4x = 16$  or  $x + 3 = 7$

3 a 6

b 9

c 7

d 3

e 1

f 5

g 4

h 2

i 8

#### Exercise 15B

1 a  $x = 2$

b  $x = 13$

c  $x = 13$

d  $y = 6$

e  $x = 9$

f  $x = 5$

g  $t = 18$

h  $y = 24$

i  $k = 18$

j  $h = 40$

2 2

3 27p

#### Exercise 15C

1 a  $x = 56$

b  $y = 2$

c  $x = 6$

d  $y = 3$

e  $w = 21$

f  $x = 72$

g  $m = 56$

h  $x = 0$

i  $h = -7$

j  $f = 60$

2 a  $-4$

b 15

3 Any valid equation such as

$3x + 8 = 2$ ,  $6x + 20 = 8$

4 a Betsy

b Second line: Amanda subtracts 1 instead of adding 1; fourth line: Amanda subtracts 2 instead of dividing by 2.

#### Exercise 15D

1 a  $k = 5$

b  $h = 28$

c  $w = 5$

d  $x = 35$

e  $y = 33$

f  $f = 23$

2 a 33

b 48

3 a  $x = 2$

b  $x = 7$

c  $y = 2\frac{4}{5}$

d  $y = 1$

e  $x = 11\frac{1}{2}$

f  $t = \frac{1}{5}$

4 Any valid equations

5 a Anwar

b First line: Brody adds 4 instead of multiplying by 5.

Second line: Brody adds 5 instead of multiplying by 5.

Fourth line: Brody subtracts 2 instead of dividing by 2.

6 a  $\frac{x + 10}{5} = 9.50$

b £37.50

7 a  $\frac{8}{3}$

b Student's own working

#### Exercise 15E

1 a  $\frac{1}{2}$

b  $1\frac{1}{5}$

c 2

d  $-2$

e  $-1$

f  $-2$

g  $-2$

h  $-1$

2 Any values that work, e.g.  $a = 2$ ,  $b = 3$  and  $c = 30$ .

3 3 cm

4 5

5 Multiplying out the brackets and collecting like terms gives  $4x - 24 = 0$  which has the solution  $x = 6$ .

6  $168^\circ$

#### Exercise 15F

1 a  $x = 2$

b  $y = 1$

c  $a = 7$

d  $t = 4$

e  $p = 2$

f  $k = -1$

g  $m = 3$

h  $s = -2$

2  $3x - 2 = 2x + 5$ ,  $x = 7$

- 3 a  $d=6$       b  $x=11$       c  $y=1$   
 d  $h=4$       e  $b=9$       f  $c=6$
- 4 a  $6x+3=6x+10$ ;  $6x-6x=10-3$ ;  $0=7$ , which is obviously false. Both sides have  $6x$ , which cancels out.  
 b Multiplying out the brackets gives  $12x+18=12x+18$  which is true for all values of  $x$ .
- 5  $8x+7+x+4=11x+5-x-4$ ,  $x=10$ , so the total value of silver coins is £1.90
- 6 a Both are expressions for the length of the rectangle so they must be equal  
 b  $70 \text{ cm}^2$
- 7 a 15  
 b  $-1$   
 c  $2(n+3)$ ,  $2(n+3)-5$   
 d  $2(n+3)-5=n$ ,  $2n+6-5=n$ ,  $2n+1=n$ ,  $n=-1$
- 8  $4x+18=3x+1+50$ ,  $x=33$ . Large bottle 1.5 litres, small bottle 1 litre

### Review questions

- 1 a  $x=6$       b  $x=9$       c  $x=\frac{2}{3}$   
 d  $x=-5$       e  $x=72$
- 2 3 years
- 3  $x=-6$
- 4 a  $x=-3$       b  $x=-3$       c  $x=-3$
- 5 c
- 6 8
- 7 Length is 5.5 m, width is 2.5 m and area is  $13.75 \text{ m}^2$ . Carpet costs £123.75
- 8 a All the sides of a square are equal  
 b  $42\frac{1}{4} \text{ m}^2$
- 9  $x=10$
- 10 14 cm
- 11 a B:  $\frac{3}{8}x$ , C:  $\frac{3}{8}x$ , D:  $\frac{1}{4}x$   
 b  $\frac{3}{8}x=300$ ,  $x=800$  cars  
 c:  $\frac{1}{4}x=500$ ,  $x=2000$  cars
- 12 No, as  $x+x+2+x+4+x+6=360$  gives  $x=87^\circ$  so the consecutive numbers (87, 89, 91, 93) are not even but odd

## Chapter 16 – Ratio and proportion and rates of change: Percentages and compound measures

### Exercise 16A

- 1 a  $\frac{2}{25}$       b  $\frac{7}{20}$       c  $\frac{9}{10}$
- 2 a 0.27      b 0.85      c 0.13
- 3 a  $\frac{3}{25}$       b  $\frac{2}{5}$       c  $\frac{9}{20}$
- 4 a 29%      b 60%      c 125%
- 5 a 28%      b 30%      c 95%  
 d 34%      e 27.5%      f 87.5%
- 6 a 0.6      b 0.075      c 0.76  
 d 0.3125      e 0.05      f 0.125
- 7 none
- 8 a 77%      b 39%      c 63%
- 9 27%
- 10 a 50%      b 20%      c 80%
- 11 a 87.5%, 75%, 62.5%, 50%, 25%  
 b 37.5%
- 12 a  $\frac{3}{5}$       b 0.6      c 60%      d 60%
- 13 a 63%, 83%, 39%, 62%, 77%  
 b English
- 14 10%
- 15 34%, 0.34,  $\frac{17}{50}$ ; 85%, 0.85,  $\frac{17}{20}$ ; 7.5%, 0.075,  $\frac{3}{40}$ ; 45%, 0.45,  $\frac{9}{20}$ ; 30%, 0.3,  $\frac{3}{10}$ ; 67%, 0.67,  $\frac{2}{3}$ ; 84%, 0.84,  $\frac{21}{25}$ ; 45%, 0.45,  $\frac{9}{20}$ ; 37.5%, 0.375,  $\frac{3}{8}$
- 16  $\frac{9}{10}$  or 90%

### Exercise 16B

- 1 a 0.88      b 0.3      c 0.25      d 0.08      e 1.15
- 2 a 78%      b 40%      c 75%      d 5%      e 110%
- 3 a £45      b £6.30      c 128.8 kg  
 d 1.125 kg      e 1.08 h      f 37.8 cm  
 g £0.12      h 2.94 m      i £7.60  
 j 33.88 min      k 136 kg      l £162
- 4 a 504      b 48
- 5 £2410

- 6 a 86%                                      b 215  
 7 8520  
 8 287  
 9 A total of 91% were allocated, leaving 9% or 8100 tickets for celebrities  
 10 990  
 11 Mon: 816, Tue: 833, Wed: 850, Thu: 799, Fri: 748  
 12 a £3.25            b 2.21 kg      c £562.80  
     d £6.51            e 42.93 m     f £24  
 13 480 cm<sup>3</sup> nitrogen, 120 cm<sup>3</sup> oxygen  
 14 13  
 15 £270  
 16 More in 2014 as it was 3% of a higher amount than in 2013

### Exercise 16C

- 1 a 1.1    b 1.03    c 1.2    d 1.07    e 1.12  
 2 a 0.92    b 0.85    c 0.75    d 0.91    e 0.88  
 3 a 391 kg    b 824.1 cm    c 253.5 g  
     d £143.50    e 736 m      f £30.24  
 4 a 731 m    b 83.52 g    c 360 cm  
     d 117 min    e 81.7 kg    f £37.70  
 5 448  
 6 No, as the total is £101. She will save £20.20, which is less than the £25 it would cost to join the club.  
 7 A 7% pay rise. 7% is £1925 per year; £150 per month is £ 1800 per year.  
 8 a £6.13 (£6.125)  
     b  $x \times 0.025$   
     c  $y \div 1.175 \times 1.2$   
 9 Offer B is the better offer:  
     Offer A gives 360 grams for £1.40, i.e. 0.389 pence per gram.  
     Offer B gives 300 grams for £1.12, i.e 0.373 pence per gram.  
     Or: Offer A is 360 for 1.40 = 2.6 g/p, offer B is 300 for 1.12 = 2.7 g/p.  
 10 c Both the same as  $1.05 \times 1.03 = 1.03 \times 1.05$   
 11 a Shop A as  $1.04 \times 1.04 = 1.0816$ , so an 8.16% increase.  
 12 £425.25  
 13  $1.10 \times 0.9 = 0.99$  (99%)

### Exercise 16D

- 1 a 25%    b 60.6%    c 46.3%    d 12.5%  
     e 41.7%    f 60%      g 20.8%    h 10%  
     i 1.9%    j 8.3%      k 45.5%    l 10.5%  
 2 32%  
 3 6.5%  
 4 33.7%  
 5 a 49.2%    b 64.5%    c 10.6%  
 6 4.9%  
 7 90.5%  
 8 Stacey had the greater percentage increase.  
     Stacey:  $(20 - 14) \times 100 \div 14 = 42.9\%$   
     Calum:  $(17 - 12) \times 100 \div 12 = 41.7\%$   
 9 Yes, as 38 out of 46 is over 80% (82.6%)  
 10 Let  $z = 100$ .  $y = 75$ ,  $x = 0.6 \times 75 = 45$ , so  $x$  is 45% of  $z$   
 11 Let  $z$  be 100,  $x = 60$ . If  $x$  is 75% of  $y$ ,  $y = 80$ , so  $y$  is 80% of  $z$ .  
 12 30% of 4800 = 1440.  $1.2 \times 4800 = 5760$ . 70% of 5760 = 4032.  $4032 - 1440 = 2592$ .  
      $2592 \div 1440 = 1.8$ , so the increase in numbers owning a mobile phone is 180%.  
 13  $31 \div 26 = 1.19$  which is a 19% increase. 31% is 5% more of the total votes cast than 26%

### Exercise 16E

- 1 a £260    b £307.50    c £323.75    d £289  
 2 a £7.50    b £9.05      c £5.80      d £10.75  
 3 a 38 h    b  $41\frac{1}{2}$  h    c 35 h      d 40 h  
 4 a Fewer hours                      b More pay per week  
 5 a £540                                      b £702  
 6 £6.90  
 7  $375 - 330 = 45$ ,  $45 \div 6 = £7.50$ .  
      $(375 - 12 \times 7.50) \div 7.50 = 38$  hours  
 8  $£1\frac{1}{4}x$  or  $£\frac{5x}{4}$   
 9 Pay is £442.50 tax is £88.50, NI is 442.50 – 88.50 – 327.45 = 26.55,  $26.55 \div 442.5 = 0.06$ , so the NI rate is 6%  
 10 407 factorises to  $1 \times 407$  or  $11 \times 37$ , so Antoine works 37 hours a week at £11 per hour.

**Exercise 16F**

- 1  $0.75 \text{ g/cm}^3$
  - 2 4 Pa
  - 3  $8.33 \text{ g/cm}^3$
  - 4 2.5 N
  - 5 32 g
  - 6  $5 \text{ m}^2$
  - 7  $120 \text{ cm}^3$
  - 8 156.8 g
  - 9 The 30 cm  $\times$  20 cm face
  - 10 The handles exert greater pressure as there is less area in contact with your hands.
  - 11 To reduce the pressure they exert, which helps prevent them sinking into soft sand.
  - 12 a 19.3 kg  
b 19.3 kg. The scale shows the mass, which is the same  
c (a) 9650 Pa, (b) 38 600 Pa
  - 13 The first statue is a fake
  - 14 The 400 g piece has a bigger volume by  $1 \text{ cm}^3$
  - 15 If the soles of the shoes and slippers are the same size he will exert the same pressure
  - 16 They are smooth and narrow so cut through snow. They have a big area so they don't push into the snow
  - 17 Lay down and slide. Less pressure is exerted on the ice
  - 18 Each wheel takes some of the mass so the pressure on the road or a weak bridge is less.
  - 19 A, D, C, B. If they have the same mass, the one with the largest area will exert the least pressure.
  - 20 a Same, as the area is the same  
b Cone, as all the mass is concentrated on the point which has negligible area.
  - 21  $0.34 \text{ m}^3$
  - 22  $1 \text{ m} \times 0.8 \text{ m} \times 0.5 \text{ m}$
  - 23 a True      b False      c False
- 3 £317.55
  - 4 £8.90
  - 5 France. Their speed limit is equivalent to 75 mph.
  - 6 a  $1.73 \div 0.04 = 43.25$  so 43 horses,  $2.64 \div 0.065 = 40.61$  so 40 cattle and  $0.95 \div 0.01 = 95$ . Total  $43 + 40 + 95 = 178$  animals.  
b Horses in field A = 43, Sheep in field B =  $2.64 \div 0.01 = 264$ , Cattle in field C,  $0.95 \div 0.065 = 14.61$ , so 14 cattle. Total  $43 + 264 + 14 = 321$  animals
  - 7 25%
  - 8 £8.98
  - 9  $8 \text{ g/cm}^3$
  - 10 0.88
  - 11 240 Pa
  - 12 Shop B
  - 13 a £9.00  
b £13.20
  - 14 £322
  - 15 Yes, the council exceeded the promise which was for a maximum of 1200 unemployed people.
  - 16 The 20% reduction will reduce the prices by more than the original VAT because they will be reducing prices by 20% of a bigger amount.
  - 17 a 150 men, 100 women    b 12%
  - 18 13%

## Chapter 17 – Ratio and proportion and rates of change: Percentages and variation

**Exercise 17A**

- 1 a £400    b £90    c £480    d £12.80
- 2 a 10.5 g    b 11.03 g    c 12.16 g    d 14.07 g
- 3 a 2550    b 2168    c 1331
- 4 a £6800    b £5440    c £3481.60
- 5 a i 51 980    ii 84 752    iii 138 886
- 6 a 8 years    b 12 years
- 7 12 years

**Review questions**

- 1 £144.00
- 2 £208.25

- 8 a £14 272.27                      b 20 years  
 9 a i 1.9 million litres  
     ii 1.6 million litres  
     iii 1.2 million litres  
     b 10th August  
 10 a 21 years                          b 21 years

**Exercise 17B**

- 1 a 800 g            b 250 m            c 60 cm  
   d £3075          e £200            f £400  
 2 80  
 3 T-shirt £8.40, Tights £1.20, Shorts £5.20,  
   Sweater £10.75, Trainers £24.80, Boots £32.40  
 4 £833.33  
 5 £300  
 6 240  
 7 £350  
 8 4750 blue bottles  
 9 £22  
 10 a £1600  
     b Yes. With 10% cut each year he earns  
        $£1440 \times 12 + £1296 \times 12 = £17\,280 + £15\,552 = £32\,832$ . With an immediate 14%  
       cut he earns  $£1376 \times 24 = £33\,024$ , so he  
       made the correct decision.

**Exercise 17C**

- 1 a £3.40            b 17                      c 12  
 2 a 100 cm                      b 20 inches  
   c 82 inches is longer  
 3 a 150                          b 6  
 4 a 22.5                        b 12  
 5 a 175 miles                b 8 hours  
 6 a £66.50                      b 175 kg  
 7 a 44                            b 84 m<sup>2</sup>  
 8 a 50  
 9 17 minutes 30 seconds  
 10 a i = 4.55G    b No, 6 gallons is 27.3 litres  
     c £5.55            d 11.9 miles per litre

**Exercise 17D**

- 1 a 12.5 watts                      b 2.5 m  
 2 8 hours

- 3 a 5 cm                              b 10 cm  
 4 a 625 cm<sup>3</sup>                      b 0.625 bar  
 5 a 6                                      b 12.5  
 6 a 27 min 31.2 sec              b 34.4 km/h  
 7 a £150                              b 36  
 8 a 3 °C                                b 12 km  
 9

x	8	24	16
y	1	$\frac{1}{3}$	$\frac{1}{2}$

- 10 a 112.5 miles                      b 7.5 miles

**Review questions**

- 1 £2120  
 2 £19 728  
 3 £112.50  
 4 a 37p                                      b 3 hours  
   c 20 minutes                            d 36 kg  
   e 1 hour                                    f 16 hours  
   g Three days                            h 21 minutes  
   i 2 hours                                    j 600 p.m.  
 5 She should choose A  
 6 £8357.35  
 7 a 3 days                                  b 2 days  
 8 £8.60  
 9 £38.50  
 10 a 6    b 100  
 11 a 4    b 32  
 12 £375  
 13 a 66    b 1.75  
 14 a 8    b 0.5  
 15

x	25	50	400
y	10	20	160

- 16 a 125                                      b  $\frac{1}{2}$   
 17  $P = k_1Q$ ,  $k_1 = \frac{P}{Q} = \frac{4}{10} = 0.4$ . So  $P = 0.4Q$ .  
 $QR = k_2 = 2 \times 8$ . So  $QR = 16$ . When  $R = 0.8$ ,  
 then  $Q = 16 \div 0.8 = 20$ .  
 So  $P = 0.4Q = 0.4 \times 20 = 8$

## Chapter 18 – ~~More complex~~ statistics

### Exercise 18A

- secondary data
  - primary data
  - primary or secondary data
  - primary or secondary data
  - primary data
  - primary data
- Plan the data collection: Choose a random sample of 30 boys and 30 girls from Year 11.

Collect the data: Ask each student to spell the same 10 words. This will avoid bias. Pick words that are often misspelt, eg accommodation, necessary, separate

Choose the best way to process and represent the data: Calculate the mean number and range for the number of correct spellings for the boys and for the girls. Draw a dual bar chart to illustrate the data.

Interpret the data and make conclusions: Compare the mean and range to arrive at a conclusion. Is there a clear conclusion or do you need to change any of the 10 words or take a larger sample?

- Plan the data collection: Choose a random sample of 20 boys and 20 girls from Year 11.

Collect the data: Ask each student, on average, how many hours of sport they play and how many hours of TV they watch each week.

Choose the best way to process and represent the data: Calculate the mean number of hours for the number of hours playing sport and the number of hours watching TV. Draw dual bar chart illustrate the data.




Interpret the data and make conclusions: Compare the means and arrive at a conclusion. Is there a clear conclusion or do you need to take a larger sample?

- An example might be:
  - 20 to each of the 6 groups
  - For example, get a list of the names of the students in alphabetical order for each group. Then choose a random sample for each one by picking every 5th student on the list.

- Decide on a sample size. A suitable sample size here is 100. As the ratio of males to females is 3:1, take a random sample in the same ratio. The sample therefore should be 75 males and 25 females.
- There are more students in Year 12 and a different number of boys and girls in both years.

	Male	Female	Total
Full time	13	7	20
Part time	4	6	10
Total number of employees			30

### Exercise 18B


- 
  - 
  - 
- Pie charts with the following angles:
  - $36^\circ, 90^\circ, 126^\circ, 81^\circ, 27^\circ$
  - $90^\circ, 108^\circ, 60^\circ, 78^\circ, 24^\circ$
  - $168^\circ, 52^\circ, 100^\circ, 40^\circ$
- Pictogram with suitable key
  - Bar chart correctly labelled
  - Vertical line chart correctly labelled
  - Pie chart with these angles:  $60^\circ, 165^\circ, 45^\circ, 15^\circ, 75^\circ$  and correctly labelled
  - Vertical line chart. It shows the frequencies, the easiest one to draw and comparisons can be made.
- 36
  - Pie charts with these angles:  $50^\circ, 50^\circ, 80^\circ, 60^\circ, 60^\circ, 40^\circ, 20^\circ$  and correctly labelled
  - Bar chart correctly labelled
  - Bar chart, because it is easier to make comparisons
- Pie charts with these angles:  $124^\circ, 132^\circ, 76^\circ, 28^\circ$  and correctly labelled
  - Split of parties can be seen at a glance
- Pie charts with these angles:  
Strings:  $36^\circ, 118^\circ, 126^\circ, 72^\circ, 8^\circ$   
Brass:  $82^\circ, 118^\circ, 98^\circ, 39^\circ, 23^\circ$
  - Overall, the strings candidates did better, as a smaller proportion scored the lowest grade. a higher proportion of brass candidates scored the highest grade.
- The missing angle is  $40^\circ$ , so  $P(\text{Don't know}) = \frac{40}{360} = \frac{1}{9}$

- 8 a  $55^\circ$     b 22    c  $33\frac{1}{3}\%$

**Exercise 18C**

- 1 a i Good positive correlation  
 ii Time increases with the number of press-ups  
 b i Strong negative correlation  
 ii You complete a crossword quicker as you get older  
 c i No correlation  
 ii Speed of cars on M1 is not related to the temperature  
 d i Weak positive correlation  
 ii Older people generally have more money saved in the bank


- 2 a and b



c about 19 cm/s

d about 34 cm

- 3 a and b




c Greta

d about 70


e about 70

- 4 a




- b Yes, as good positive correlation

- 5 a



- b Little correlation, so cannot draw a line of best fit

- 6 a and b



c about 2.4 km

d about 8 minutes

e you cannot extrapolate values from a scatter diagram or the data may change for longer journeys

- 7 about 23 mph

- 8 Points showing a line of best fit sloping down from top left to bottom right



**Exercise 18D**

- 1 a i  $30 < x \leq 40$  ii 29.5 iii  $30 < x \leq 40$   
 b i  $0 < y \leq 100$  ii 158.3 iii  $100 < y \leq 200$   
 c i  $5 < z \leq 10$  ii 9.43 iii  $5 < z \leq 10$   
 d i 7–9 ii 8.41 iii 7–9
- 2 a  $100 < m \leq 120$  b 10.86 kg c 108.6 g
- 3 a  $175 < h \leq 200$   
 b 31%  
 c 193.3 hours  
 d No, the mean was under 200.
- 4 a Yes, average distance is 11.7 miles per day.  
 b Because shorter runs will be done faster which will affect the average  
 c Yes, because the shortest could be 1 mile, the longest 25 miles.
- 5 24
- 6 Soundbuy; average increases are Soundbuy 17.7p, Springfields 18.7p, Setco 18.2p
- 7 a 160 b 52.6 minutes  
 c Modal group d 65%
- 8 The first 5 and the 10 are the wrong way round.
- 9 Find the midpoint of each group, multiply that by the frequency and add those products. Divide that total by the total frequency.

**Review questions**

- 1 a 18 b  $80^\circ$
- 2 a i Diagram C ii Diagram A iii Diagram B  
 b Diagram A: strong negative correlation, diagram B: no correlation, diagram C: strong positive correlation
- 3 a/b Time on horizontal axis from 0 to 20 and Distance (km) on vertical axis from 0 to 10 with the following points plotted: (3, 1.7) (17, 8.3) (11, 5.1) (13, 6.7) (9, 4.7) (15, 7.3) (8, 3.8) (11, 5.7) (16, 8.7) (10, 5.3) and with line of best fit drawn.  
 c/d Answers depend on student's plotting
- 4 a  $10 < t \leq 20$  b  $10 < t \leq 20$  c 19 minutes
- 5 a Because over half the students have more than £10 pocket money, so the mean must be more than £10  
 b £11.00
- 6 17.15 mph




**Chapter 19 – Geometry and measures: Constructions and loci****Exercise 19A**

- 1 a  $BC = 2.9$  cm,  $\angle B = 53^\circ$ ,  $\angle C = 92^\circ$   
 b  $EF = 7.4$  cm,  $ED = 6.8$  cm,  $\angle E = 50^\circ$   
 c  $\angle G = 105^\circ$ ,  $\angle H = 29^\circ$ ,  $\angle I = 46^\circ$   
 d  $\angle J = 48^\circ$ ,  $\angle L = 32^\circ$ ,  $JK = 4.3$  cm  
 e  $\angle N = 55^\circ$ ,  $ON = OM = 7.0$  cm  
 f  $\angle P = 51^\circ$ ,  $\angle R = 39^\circ$ ,  $QP = 5.7$  cm
- 2 a Students can check one another's triangles.  
 b  $\angle ABC = 44^\circ$ ,  $\angle BCA = 79^\circ$ ,  $\angle CAB = 57^\circ$
- 3 a 5.9 cm  
 b  $18.8$  cm<sup>2</sup>
- 4  $BC = 2.6$  cm, 7.8 cm
- 5 a 4.3  
 b  $10.8$  cm<sup>2</sup>
- 6 a 4.3 cm  
 b  $34.4$  cm<sup>2</sup>
- 7 An equilateral triangle of side 4 cm
- 8 Even with all three angles, you need to know at least one length


**Exercise 19B**

- 1–9 Practical work; check students' constructions
- 10 The centre of the circle
- 11 Start with a base line AB; then construct a perpendicular to the line from point A. At point B, construct an angle of  $60^\circ$ . Ensure that the line for this  $60^\circ$  angle crosses the perpendicular line; where they meet will be the final point C.
- 12–14 Practical work; check students' constructions



**Exercise 19C**

- 1 Circle with radius:  
 a 2 cm b 4 cm c 5 cm
- 2 a  b  c 

3 a Circle with radius 4 m





b



6








7 Construct the bisector of angle BAC and the perpendicular bisector of the line AC.



9 Start with a base line, AB, 3cm long. At point A, draw a few points all 3cm away from a towards the upper right side. Lightly join these dots with an arc. You can now find the point that is 3cm away from point B and draw the equilateral triangle.

10 Gary is correct about the triangle inside, but not a triangle outside as there will be three straight lines, parallel to each side of the triangle, then these straight lines will be joined with arcs centred on the vertex between the lines.

**Exercise 19D**



7 a Sketch should show a circle of radius 6 cm around London and one of radius 4 cm around Glasgow.

b No  
c Yes

8 a Yes  
b Sketch should show a circle of radius 4 cm around Leeds and one of radius 4 cm around Exeter. The area where they overlap should be shaded.

9 a This is the perpendicular bisector of the line from York to Birmingham. It should pass just below Manchester and just through the top of Norwich.

b Sketch should show a circle of radius 7 cm around Glasgow and one of radius 5 cm around London.

c The transmitter can be built anywhere on line constructed in part a that is within the area where the circles shown in part b overlap.

10 Sketch should show two circles around Birmingham, one of radius 3 cm and one of radius 5 cm. The area of good reception is the area between the two circles.

**11** Sketch should show a circle of radius 6 cm around Glasgow, two circles around York, one of radius 4 cm and one of radius 6 cm and a circle around London of radius 8 cm. The small area in the Irish Sea that is between the two circles around York and inside both the circle around Glasgow and the circle around London is where the boat can be.

**12** Sketch should show two circles around Newcastle upon Tyne, one of radius 4 cm and one of radius 6 cm, and two circles around Bristol, one of radius 3 cm and one of radius 5 cm. The area that is between both pairs of circles is the area that should be shaded.

**13** Sketch should show the perpendicular bisector of the line running from Newcastle upon Tyne to Manchester and that of the line running from Sheffield to Norwich. Where the lines cross is where the oil rig is located.


**14** Sketch should show the perpendicular bisector of the line running from Glasgow to Norwich and that of the line running from Norwich to Exeter. Where the lines cross is where Fred's house is.

**15** Leeds


**16** No, if you accurately draw a diagram showing the path of each boat, you will find the boat leaving from point B meets the path of the other boat in much shorter time as it's a smaller distance than from A to the cross over point.

**17** On a map, draw a straight line from Newcastle to Bristol, construct the line bisector, then the search will be anywhere on the sea along that line.

**18 a-d**




**19**




## Review questions

**1 a - b**




**c** You should have measured the error of the angle size and converted that to a percentage error.


**2**




**3**



**4**




5 a



b ..... an equal distance from a and B, or equidistant from a and B.

6




7 a Student's construction

b  $20 \text{ cm}^2$


8 a Student's construction

b  $24 \text{ cm}^2$

9 Oliver is incorrect as the correct shape is:



10



11



## Chapter 20 – Geometry and measures: Curved shapes and pyramids

### Exercise 20A

- 1 a i 5.86 cm      ii  $22.3 \text{ cm}^2$   
 b i 5.24 cm      ii  $13.1 \text{ cm}^2$   
 c i 10.99 cm     ii  $19.24 \text{ cm}^2$   
 d i 16.34 cm     ii  $98.02 \text{ cm}^2$

- 2 a i 6.8 cm      ii 18.8 cm  
 b i 7.9 cm      ii 25.9 cm  
 c i 8.4 cm      ii 16.4 cm  
 d i 17.5 cm     ii 37.5 cm  
 e i 3.3 cm      ii 20.3 cm

3 Arc length =  $2\pi$ , area of sector =  $6\pi$

4 a  $270^\circ$       b 51.8 cm      c 73.8 cm

5 area of sector =  $\frac{1}{4} \times \pi \times 8^2 = 16\pi$ , area of circle =  $\pi \times 4^2 = 16\pi$

6 a  $28.6^\circ$       b  $66.8^\circ$       c  $54.9^\circ$

7 Area B =  $107 \text{ cm}^2$ , area B =  $173 \text{ cm}^2$ . Half of B is 86.5 which is less than 107, so Gabby is wrong, the area of a is more than half the area of B.

8 43.6 cm

9 a  $66.8^\circ$       b  $10 \text{ cm}^2$

**Exercise 20B**

- 1 a  $56 \text{ cm}^3$     b  $168 \text{ cm}^3$   
 c  $1040 \text{ cm}^3$     d  $84 \text{ cm}^3$   
 e  $160 \text{ cm}^3$
- 2  $\frac{1}{3} \text{ base area} \times h = \frac{1}{3} \times 9 \times 9 \times 10 = 270 \text{ cm}^3$
- 3 a  $73.3 \text{ m}^3$     b  $45 \text{ m}^3$     c  $3250 \text{ cm}^3$
- 4 208 g
- 5  $1.5 \text{ g/cm}^3$
- 6  $\frac{1}{3} \times 6.4 \times 6.4 \times H = 81.3$   
 so  $h = \frac{3 \times 81.3}{6.4 \times 6.4} = 5.59 = 6.0 \text{ cm}$  (2 sf)
- 7 14.4 cm

**Exercise 20C**

- 1 a  $72 \text{ cm}^2$     b  $172.5 \text{ cm}^2$
- 2 Area of base =  $9^2 = 81 \text{ cm}^2$ , area of one triangle =  $\frac{1}{2} \times 9 \times 10 = 45 \text{ cm}^2$ , total surface area =  $(4 \times 45) + 81 = 261 \text{ cm}^2$ .
- 3 Total area =  $32.46 \text{ m}^2$ , so 3 tins are needed
- 4 Surface areas are a =  $1200 \text{ cm}^2$ , b =  $576 \text{ cm}^2$ , c =  $864 \text{ cm}^2$ , d =  $526.5 \text{ cm}^2$ , Volumes are a =  $960 \text{ cm}^3$ , b =  $512 \text{ cm}^3$ , c =  $960 \text{ cm}^3$ , d =  $540 \text{ cm}^3$ . This does not show that the larger the volume, the larger the surface area, so Andrew is incorrect.
- 5 Yes, the area is  $218.7 \text{ m}^2$  and will cost £2515.05 to retille.

**Exercise 20D**

- 1 a  $17.5 \text{ cm}^2$     b  $103 \text{ cm}^2$     c  $94.2 \text{ cm}^2$
- 2 c Radii are 1.1 cm, 4.7 cm, 5 cm
- 3

Sector	Area of sector	Radius of cone, $r$	Slant height, $l$	$\pi \times r \times l$
a	$17.5 \text{ cm}^2$	1.1 cm	5 cm	$17.3 \text{ cm}^2$
b	$103 \text{ cm}^2$	4.7 cm	7 cm	$103 \text{ cm}^2$
c	$94.2 \text{ cm}^2$	5 cm	6 cm	$94.2 \text{ cm}^2$

- 4 a They are the same  
 b Curved surface area =  $\pi r l$

**Exercise 20E**

- 1 a i  $151 \text{ cm}^3$     ii  $176 \text{ cm}^2$   
 b i  $75 \text{ cm}^3$     ii  $108 \text{ cm}^2$   
 c i  $872 \text{ cm}^3$     ii  $550 \text{ cm}^2$   
 d i  $3560 \text{ cm}^3$     ii  $1426 \text{ cm}^2$   
 e i  $314 \text{ cm}^3$     ii  $283 \text{ cm}^2$   
 f i  $1018 \text{ cm}^3$     ii  $679 \text{ cm}^2$
- 2 935 g
- 3  $\pi r l + \pi r^2 = 3 \pi l + 9 \pi = 75.4$   
 $3 \pi l = 75.4 - 28.27 = 47.13$   
 $l = \frac{47.13}{3\pi} = 5 \text{ cm}$
- 4 a  $816 \pi \text{ cm}^3$     b  $720 \pi \text{ mm}^3$
- 5 The total surface area, to 3sf, is  $75.4 \text{ cm}^2$ , so Mandy is correct if the figure was rounded to 1 sf. Neil is incorrect as he has cut off his final answer instead of rounding it.

**Exercise 20F**

- 1 a  $113 \text{ cm}^3$     b  $905 \text{ cm}^3$     c  $4189 \text{ cm}^3$
- 2 a  $113 \text{ cm}^2$     b  $314 \text{ cm}^2$     c  $616 \text{ cm}^2$
- 3 Volume =  $4500 \pi \text{ cm}^3$ , surface area =  $900 \pi \text{ cm}^2$
- 4 a  $1960 \text{ cm}^2$  (to 3 sf)  
 b  $7444 \text{ cm}^3$  (to nearest unit)
- 5 125 cm
- 6 6232

**Review questions**

- 1 a 43.9 cm    b 112  $\text{cm}^2$
- 2  $49\,000 \text{ m}^3$
- 3 Volume of top cube =  $30^3 = 27\,000$ , volume of bottom pyramid =  $\frac{1}{3} \times 30^2 \times 10 = 3000 \text{ cm}^3$ , total volume = top + bottom =  $27\,000 + 3\,000 = 30\,000 \text{ cm}^3$
- 4 Bottle will hold =  $\pi r^2 h = \pi \times 4.5^2 \times 25 = 1590.4 \text{ cm}^3$ . One glass will hold  $\frac{1}{3} \pi r^2 h = \frac{1}{3} \times \pi \times 2.5^2 \times 6 = 39.27 \text{ cm}^3$ . So  $1590.4 \div 39.27 = 40.5$  glasses can be filled. David is wrong as just over 40 glasses will be filled.
- 5 One bearing has volume =  $\frac{4}{3} \pi r^3 = \frac{4}{3} \times \pi \times 0.75^3 = 1.767 \text{ cm}^3$ , so one ball bearing has a mass of  $1.767 \times 7.3 = 12.900 \text{ g}$ . 20 ball bearings have a mass of 258.00 g or 258 g (nearest whole number).
- 6 0.5 cm

- 7  $34.6 \text{ m}^2$
- 8 **a** A is largest at  $1018 \text{ cm}^2$   
**b** B is smallest at  $1003 \text{ cm}^2$   
 Surface area of C is  $1016 \text{ cm}^2$ .
- 9  $0.825 \text{ m}^3$
- 10  $7.25 \text{ cm}$
- 11 Neither of them is correct as the surface area of the sphere is  $1257 \text{ cm}^2$ , surface area of the cube is  $1262 \text{ cm}^2$  but the surface area of the cone is  $1282 \text{ cm}^2$ , so the cone has the greatest surface area
- 12  $91.5 \text{ cm}$

## Chapter 21 – Algebra: Number and sequences

### Exercise 21A

- 1 **a**  $11111 \times 11111 = 123454321$ ,  
 $111111 \times 111111 = 12345654321$   
**b**  $99999 \times 99999 = 9999800001$ ,  
 $999999 \times 999999 = 999998000001$
- 2 **a**  $7 \times 8 = 7^2 + 7$ ,  $8 \times 9 = 8^2 + 8$   
**b**  $50 \times 51 = 2550$ ,  $60 \times 61 = 3660$
- 3 **a**  $1 + 2 + 3 + 4 + 5 + 4 + 3 + 2 + 1 = 25 = 5^2$ ,  
 $1 + 2 + 3 + 4 + 5 + 6 + 5 + 4 + 3 + 2 + 1 = 36 = 6^2$   
**b**  $21 + 23 + 25 + 27 + 29 = 125 = 5^3$ ,  
 $31 + 33 + 35 + 37 + 39 + 41 = 216 = 6^3$
- 4 **a**  $1 + 6 + 15 + 20 + 15 + 6 + 1 = 64$ ,  
 $1 + 7 + 21 + 35 + 35 + 21 + 7 + 1 = 128$   
**b**  $12\ 345\ 679 \times 45 = 555\ 555\ 555$ ,  
 $12\ 345\ 679 \times 54 = 666\ 666\ 666$
- 5 **a**  $1^3 + 2^3 + 3^3 + 4^3 = (1 + 2 + 3 + 4)^2 = 100$ ,  
 $1^3 + 2^3 + 3^3 + 4^3 + 5^3 = (1 + 2 + 3 + 4 + 5)^2 = 225$   
**b**  $36^2 + 37^2 + 38^2 + 39^2 + 40^2 = 41^2 + 42^2 + 43^2 + 44^2$ ,  
 $55^2 + 56^2 + 57^2 + 58^2 + 59^2 + 60^2 = 61^2 + 62^2 + 63^2 + 64^2 + 65^2$
- 6 **a**  $12345678987654321$   
**b**  $999999998000000001$   
**c**  $12^2 + 12$   
**d**  $8190$   
**e**  $81 = 9^2$   
**f**  $512 = 8^3$   
**g**  $512$   
**h**  $999\ 999\ 999$   
**i**  $(1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9)^2 = 2025$

- 7  $1 + 500 = 501$ ,  $2 + 499 = 501$ , ...,  $250 + 251 = 501$ ,  $250 \times 501 = 125250$

### Exercise 21B

- 1 **a** 21, 34: add previous 2 terms  
**b** 49, 64: next square number  
**c** 47, 76: add previous 2 terms
- 2 15, 21, 28, 36
- 3 61, 91, 127
- 4 6, 11, 16, 21, 26 ...
- 5  $\frac{1}{2}, \frac{3}{5}, \frac{2}{3}, \frac{5}{7}, \frac{3}{4}$
- 6 **a** 6, 10, 15, 21, 28  
**b** It is the sums of the natural numbers, or the numbers in Pascal's triangle or the triangular numbers.
- 7 **a** 2, 6, 24, 720    **b** 69!
- 8 X. There are 351 ( $1 + 2 + \dots + 25 + 26$ ) letters from a to Z.  $3 \times 351 = 1053$ .  $1053 - 26 = 1027$ ,  $1027 - 25 = 1002$ , so, as Z and Y are eliminated, the 1000th letter must be X.
- 9 364: Daily totals are 1, 3, 6, 10, 15, 21, 28, 36, 45, 55, 66, 78 (these are the triangular numbers). Cumulative totals are: 1, 4, 10, 20, 35, 56, 84, 120, 165, 220, 286, 364.
- 10 29 and 41
- 11 No, because in the first sequence, the terms are always one less than in the 2nd sequence
- 12  $4n - 2 = 3n + 7$  rearranges as  $4n - 3n = 7 + 2$ , so  $n = 9$

### Exercise 21C

- 1 **a** 13, 15,  $2n + 1$     **b** 25, 29,  $4n + 1$   
**c** 33, 38,  $5n + 3$     **d** 32, 38,  $6n - 4$   
**e** 20, 23,  $3n + 2$     **f** 37, 44,  $7n - 5$   
**g** 21, 25,  $4n - 3$     **h** 23, 27,  $4n - 1$   
**i** 17, 20,  $3n - 1$     **j**  $-8, -18, -10n + 42$   
**k** 4, 0,  $-4n + 24$     **l**  $-1, -6, -5n + 29$
- 2 **a**  $3n + 1$ , 151    **b**  $2n + 5$ , 105  
**c**  $5n - 2$ , 248    **d**  $4n - 3$ , 197  
**e**  $8n - 6$ , 394    **f**  $n + 4$ , 54  
**g**  $5n + 1$ , 251    **h**  $8n - 5$ , 395  
**i**  $3n - 2$ , 148    **j**  $3n + 18$ , 168  
**k**  $-7n + 47$ ,  $-303$     **l**  $-8n + 41$ ,  $-359$
- 3 **a** 33rd    **b** 30th  
**c** 100th = 499




- 14 a i 4, 16, 64, 256, 1024  
 ii 7, 49, 343, 2401, 16807  
 iii 5, 25, 125, 625, 3125  
 iv 9, 81, 729, 6561, 59 049  
 v 8, 64, 512, 4096, 32 768  
 b i 60      ii 40
- 15 2. Only 2 plus a prime will give another prime, but not always. This is because after 2, all primes are odd, and odd + odd is always an even number. No even numbers after 2 are prime.
- 16 a There are many answers,  $5 + 31 = 36$ ,  $7 + 29 = 36$ ,  $2 + 47 = 49$  etc.  
 b There are many answers,  $49 - 36 = 13$ ,  $81 - 64 = 17$
- 17 a 3, 6, 11, 18, 27  
 b 102  
 c 20th  
 d No. Because the terms of the sequence  $n^2$  alternate between even and odd, and adding 2 to each term will not change this even-odd pattern.

### Exercise 21E



- b  $4n - 3$   
 c 97  
 d 50th diagram



- b  $2n + 1$       c 121      d 49th set
- 3 a 18      b  $4n + 2$       c 12
- 4 a 5 cm, 8 cm, 11 cm, 14 cm  
 b i 20 cm      ii  $(3n + 2)$  cm      iii 152 cm  
 c 332

- 5 a i 24      ii  $5n - 1$       iii 224  
 b 25
- 6 a i 20      ii 162  
 b 79.8 km
- 7 a i 14      ii  $3n + 2$       iii 41  
 b 66
- 8 a i 5      ii  $n$       iii 18  
 b 20 tins
- 9  $2n - 1$
- 10  $(n + 1)^2$
- 11 a  $2^n$   
 b i  $100 \times 2^{n-1}$  ml      ii 1600 ml  
 c Next sizes after super giant are 3.2 l, 6.4 l and 12.8 l with weights of 3.2 kg, 6.4 kg and 12.8 kg, so the largest size is 6.4 litres.
- 12 The  $n$ th term is  $\left(\frac{3}{4}\right)^n$ , so as  $n$  gets very large, the unshaded area gets smaller and smaller and eventually it will be zero; so the shaded area will eventually cover the triangle.
- 13 Yes, as the number of matches is 12, 21, 30, 39, ... which is  $9n + 3$ ; so he will need  $9 \times 20 + 3 = 183$  matches for the 20th step and he has  $5 \times 42 = 210$  matches.
- 14 a 20      b 120
- 15 Alex's answer gives  $4(n + 2) = 4n + 8$   
 Colin's method gives  $4n + 4$   
 Ed's method gives  $4(n + 1) = 4n + 4$   
 Gail's method gives  $2 \times n + 2(n + 2) = 2n + 2n + 4 = 4n + 4$   
 Linear sequence is 8, 12, 16, 20, .... Which has an  $n$ th term of  $4n + 4$  so they are all valid methods except for Alex who forgot that the corners overlap and should have taken the 4 overlapping corners away to get  $4n + 8 - 4 = 4n + 4$
- 16 a £128      b £31      c £2048

### Review questions

- 1 No. Sequence is 7, 10, 13, 16, 19, 22, 25, 28, ... so the first 3 odd terms are prime but 25 is not prime.
- 2 a 13, 9  
 b Subtract 4  
 c All the terms are odd and  $-4$  is even



- 3 a i 40, 34 ii 32, 64  
b 4, 16, 64
- 4 a Always even b Odd or even
- 5 0, 3, 8
- 6 a 5, 3  
b 15, 14.5  
c He is right as when it gets nearly to 14,  
the next term will be  $\frac{14 + 14}{2} = 14$
- 7 a i 6, 11, 16 ii All terms end in 1 or 6  
b No. When  $n = 1$  then  $(1 + 2)^2 - 9 = 9 - 9 = 0$  and after that  $(n + 2)^2$  is always greater than 9
- 8 a  $4n + 1$   
b All the terms are odd and 112 is even  
c 28<sup>th</sup> term is 113
- 9  $n$ th term is  $5n + 1$ .  $5 \times 150 + 1 = 751$
- 10 a  $6n + 3$   
b No,  $3n + 2$  generates the sequence 5, 8, 11, 14, 17, 20, 23, ... so the even terms of this sequence are always 1 less than the terms of the original sequence
- 11 a Always odd b Always odd
- 12 For example:  $0.5^2 = 0.25$ ,  $0.25 < 0.5$
- 13 a  $2 \times n$  is always even, any even number plus 1 is odd  
b Odd  $\times$  odd = odd, even  $\times$  even = even

## Chapter 22 – Geometry and measures: Right-angled triangles

### Exercise 22A

- 1 b  $90^\circ$  c 9, 16, 25 d  $9 + 16 = 25$
- 2 b  $90^\circ$  c 25, 144, 169 d  $25 + 144 = 169$
- 3 b  $90^\circ$  c 36, 64, 100 d  $36 + 64 = 100$
- 4  $c^2 = a^2 + b^2$
- 5 b 7.8 cm c 7.8 cm

### Exercise 22B

- 1 a 10.3 cm b 5.9 cm c 8.5 cm  
d 20.6 cm e 18.6 cm f 17.5 cm  
g 32.2 cm h 2.4 m i 500 m  
j 5 cm k 13 cm l 10 cm
- 2 14.4 cm
- 3 Diagonal<sup>2</sup> =  $6^2 + 6^2 = 72$ , diagonal =  $\sqrt{72} = 8.485284 = 8.5$  cm to 1 decimal place

- 4 50 cm, 1.2 m and 1.3 m or 1.5 m, 2 m and 2.5 m
- 5 Because  $6^2 + 7^2 = 85$  and  $\sqrt{85} = 9.2$  not 10
- 6 a 15 b 13 c 7.3 d 14.2
- 7 a 9, 12, 15  
b 25, 60, 65  
c Any six more Pythagorean triples, for example 10, 24, 26

### Exercise 22C

- 1 a 15.0 cm b 14.7 cm c 6.3 cm  
d 18.3 cm e 5.4 cm f 217.9 m  
g 0.4 cm h 8.0 m
- 2 a 20.8 m b 15.5 cm c 15.5 m  
d 12.4 cm e 22.9 m f 19.8 m  
g 7.1 m h 0.6 m
- 3 a 5.0 m b 6.0 m c 3.0 m d 50.0 cm
- 4 The  $12 \times 30$  has diagonal length of 32.31 cm, the  $11 \times 30$  has a diagonal length of 31.95 cm, so the  $11 \times 30$  is the closest to 32cm.
- 5 Many different combinations are possible, such as: 8 cm and 11.5 cm, 10 cm and 9.8 cm, 12 cm and 7.2 cm.
- 6  $80^2 + 60^2 = 6400 + 3600 = 10\,000 = 100^2$
- 7 a Student's own diagram  
b i The large square is 17 by 17 giving 289 square units.  
ii The red and yellow triangles all have shorter lengths of 5 and 12, with an area of 30 square units each. The four red triangles have an area of  $4 \times 30 = 120$  square units  
iii The area of the inner square (green and yellow) must be  $289 - 4 \times 30 = 169$ , so the hypotenuse of the red and yellow triangles must be  $\sqrt{169} = 13$
- c The above working shows that  $5^2 + 12^2 = 13^2$

### Exercise 22D

- 1 No. The foot of the ladder is about 6.6 m from the wall.
- 2 2.06 m
- 3 11.3 m
- 4 About 17 minutes
- 5 27.4 m
- 6 4.58 m
- 7 a 3.87 m b 1.74 m


- 8 3.16 m
- 9 This creates a right-angled triangle with two short sides of 5 and 12. Use Pythagoras' theorem to show length of line  $= \sqrt{(5 + 12^2)} = 13$
- 10 a 4.85 m  
b 4.83 m (There is only a small difference.)
- 11 Yes, because  $24^2 + 7^2 = 25^2$
- 12 6 cm
- 13 He is partly correct. The perimeter must be larger than 20 cm or the rectangle has no width, and the area is largest when it's a square, giving a perimeter of 28.3 cm (3sf). So he should have said the perimeter is between 20 and 28.3 cm.

**Exercise 22E**

- 1 a 32.2 cm<sup>2</sup> b 2.83 cm<sup>2</sup> c 50.0 cm<sup>2</sup>

- 2 22.2 cm<sup>2</sup>

- 3 15.6 cm<sup>2</sup>



- b The areas are 12 cm<sup>2</sup> and 13.6 cm<sup>2</sup> respectively, so triangle with 6 cm, 6 cm, 5 cm sides has the greater area.

- 5 a  b 166.3 cm<sup>2</sup>

- 6 259.8 cm<sup>2</sup>

c i


Triangle	Opposite	Adjacent	Hypotenuse	Opposite Hypotenuse	Adjacent Hypotenuse	Opposite Adjacent
A	1.3	2.3	2.6	0.5	0.88	0.57
B	1.1	2	2.2	0.5	0.91	0.55
C	1	1.7	2	0.5	0.85	0.59
D	0.7	1.3	1.4	0.5	0.93	0.54
E	1	1.7	2	0.5	0.85	0.59
F	1.2	2	2.3	0.5	0.87	0.6

ii Regardless of the size of the triangle the ratios are similar in each case.


- 7 a No, areas vary from 24.5 cm<sup>2</sup> to 27.7 cm<sup>2</sup>  
b No, the equilateral triangle gives the largest area.  
c The closer the isosceles triangle gets to an equilateral triangle the larger its area becomes.
- 8 The right-angled triangle made with hypotenuse 6.5 m and base  $7.4 \div 2 = 3.7$  gives the height of the triangle as 5.344 cm. Use area  $= \frac{1}{2} \times 7.4 \times 5.344$  to give 19.7733 which rounds to 19.8 m<sup>2</sup> (3sf)
- 9 48 cm<sup>2</sup>
- 10 a 10 cm b 26 cm c 9.6 cm
- 11 6 or 8 cm
- 12 Andrew didn't round off any answers until the last calculation, and Ollly used a rounded off value to find an intermediate result.

**Exercise 22F**

- 1 a For example:



- b For example:




ii Regardless of the size of the triangle the ratios are similar in each case.

- 2 **b** Yes, the pattern is the same.
- 3 **a** The answer the calculator gives is almost the same as the  $\frac{\text{opposite}}{\text{hypotenuse}}$ .
- b** The answer the calculator gives is almost the same as the  $\frac{\text{adjacent}}{\text{hypotenuse}}$ .
- c** The answer the calculator gives is almost the same as the  $\frac{\text{opposite}}{\text{adjacent}}$ .
- d** Yes
- e** Match up calculator result with student table result from above.

### Exercise 22G

- 1 **a**  $x$  is opp,  $y$  is hyp,  $w$  is adj  
**b**  $x$  is adj,  $y$  is opp,  $w$  is hyp  
**c**  $x$  is hyp,  $y$  is opp,  $w$  is adj  
**d**  $x$  is adj,  $y$  is hyp,  $w$  is opp  
**e**  $x$  is hyp,  $y$  is opp,  $w$  is adj  
**f**  $x$  is adj,  $y$  is hyp,  $w$  is opp
- 2 **b**  $\sin \theta = \frac{5}{13}$ ,  $\cos \theta = \frac{12}{13}$ ,  $\tan \theta = \frac{5}{12}$   
**c**  $\sin \theta = \frac{9}{41}$ ,  $\cos \theta = \frac{40}{41}$ ,  $\tan \theta = \frac{9}{40}$
- 3 **a**  $\sin \theta = \frac{12}{20}$ ,  $\cos \theta = \frac{16}{20}$ ,  $\tan \theta = \frac{12}{16}$   
**b**  $\sin \theta = \frac{5}{13}$ ,  $\cos \theta = \frac{12}{13}$ ,  $\tan \theta = \frac{5}{12}$   
**c**  $\sin \theta = \frac{24}{26}$ ,  $\cos \theta = \frac{10}{26}$ ,  $\tan \theta = \frac{24}{10}$   
**d**  $\sin \theta = \frac{20}{25}$ ,  $\cos \theta = \frac{15}{25}$ ,  $\tan \theta = \frac{20}{15}$   
**e**  $\sin \theta = \frac{36}{39}$ ,  $\cos \theta = \frac{15}{39}$ ,  $\tan \theta = \frac{36}{15}$   
**f**  $\sin \theta = \frac{40}{50}$ ,  $\cos \theta = \frac{30}{50}$ ,  $\tan \theta = \frac{40}{30}$
- 4 **b**  $\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{5}{7}$     **c**  $\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{7}{10}$   
**d**  $\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{8}{11}$     **e**  $\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{7}{12}$   
**f**  $\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{5}{13}$
- 5 If the triangle is right-angled, then the sides are 3, 4 and 5. This is a Pythagorean triple and so he could be correct.




- 7 **a** **i**  $\tan x = \frac{3}{4}$     **ii**  $\sin x = \frac{3}{5}$     **iii**  $\cos x = \frac{4}{5}$   
**b** **i**  $\tan y = \frac{4}{3}$     **ii**  $\sin y = \frac{4}{5}$     **iii**  $\cos y = \frac{3}{5}$   
**c** **i** the same    **ii** the same  
**d** One is the reciprocal of the other

### Exercise 22H

- 1 **a** 0.682    **b** 0.829    **c** 0.922  
**d** 1    **e** 0.707    **f** 0.342  
**g** 0.375    **h** 0
- 2 **a** 0.731    **b** 0.559    **c** 0.388  
**d** 0    **e** 0.707    **f** 0.940  
**g** 0.927    **h** 1
- 3  $45^\circ$
- 4 **a** **i** 0.574    **ii** 0.574  
**b** **i** 0.208    **ii** 0.208  
**c** **i** 0.391    **ii** 0.391  
**d** Same  
**e** **i**  $\sin 15^\circ$  is the same as  $\cos 75^\circ$   
**ii**  $\cos 82^\circ$  is the same as  $\sin 8^\circ$   
**iii**  $\sin x$  is the same as  $\cos (90^\circ - x)$
- 5 **a** 0.933    **b** 1.48    **c** 2.38  
**d** Infinite    **e** 1    **f** 0.364  
**g** 0.404    **h** 0
- 6 **a** 0.956    **b** 0.899    **c** 2.16  
**d** 0.999    **e** 0.819    **f** 0.577  
**g** 0.469    **h** 0.996
- 7 Tan has some values  $> 1$
- 8 **a** 10.7    **b** 5.40  
**c** Infinite    **d** 0
- 9 **a** 3.56    **b** 8.96  
**c** 28.4    **d** 8.91
- 10  $(4.5315389 + 4.458869) - (6 + 0) = 8.9904079 - 6 = 2.990408 = 3.0$  (2 sf)
- 11 0.6

### Exercise 22I


- 1 a 9.5cm                      b 3.8cm  
 c 4.4cm                      d 8.3 cm  
 e 12.0cm                    f 4.2cm  
 g 10.3cm                    h 19.6cm  
 i 5.1cm                      j 9.2cm  
 k 6.1cm                      l 13.3cm
- 2 Yes because one side =  $10 \sin 60^\circ = 8.7$  cm (1 dp) and the other  $10 \cos 60^\circ = 5$  cm
- 3 Opposite =  $8 \tan 38^\circ = 6.25$  cm (2 dp)
- 4  $\sin x = \frac{\text{opp}}{\text{hyp}} = \frac{P}{Q}$ ,  $\cos y = \frac{\text{adj}}{\text{hyp}} = \frac{P}{Q}$  so where  $x$  and  $y$  are the two acute angles in a right-angled triangle, the sine of one is equal to the cosine of the other.




### Exercise 22J


- 1 a  $30.0^\circ$                     b  $51.7^\circ$                     c  $39.8^\circ$   
 d  $61.3^\circ$                     e  $87.4^\circ$                     f  $45.0^\circ$
- 2 a  $60.0^\circ$                     b  $50.2^\circ$                     c  $2.6^\circ$   
 d  $45.0^\circ$                     e  $78.5^\circ$                     f  $45.6^\circ$
- 3 a  $31.0^\circ$                     b  $20.8^\circ$                     c  $41.8^\circ$   
 d  $46.4^\circ$                     e  $69.5^\circ$                     f  $77.1^\circ$
- 4 a  $53.1^\circ$                     b  $41.8^\circ$                     c  $44.4^\circ$   
 d  $56.4^\circ$                     e  $2.4^\circ$                     f  $22.6^\circ$
- 5 a  $36.9^\circ$                     b  $48.2^\circ$                     c  $45.6^\circ$   
 d  $33.6^\circ$                     e  $87.6^\circ$                     f  $67.4^\circ$
- 6 a  $31.0^\circ$                     b  $37.9^\circ$                     c  $15.9^\circ$   
 d  $60.9^\circ$                     e  $57.5^\circ$                     f  $50.2^\circ$
- 7 a His statement is true.  
 b It is true for cosine for the same reason as sine. It is not true for tangent because  $\tan x$  is larger than 1 when the opposite is larger than the adjacent.  
 c Largest value 1
- 8 a i  $17.5^\circ$                     ii  $72.5^\circ$                     iii  $90^\circ$   
 b Yes
- 9 a 61.0                    b 26.4                    c 36.9                    d 45.6  
 e 43.3                    f 54.3                    g 51.3                    h 49.4  
 i 36.9                    j 41.4                    k 39.5                    l 69.4

### Exercise 22K

- 1 
- a The opposite side is  $\sqrt{3}$  from Pythagoras.  $\tan 60^\circ = \frac{O}{A} = \frac{\sqrt{3}}{1} = \sqrt{3}$
- b  $\sin 60^\circ = \frac{O}{H} = \frac{\sqrt{3}}{2}$



- 2 a The adjacent side is  $\sqrt{3}$  from Pythagoras.  $\tan 30^\circ = \frac{O}{A} = \frac{1}{\sqrt{3}}$
- b  $\cos 30^\circ = \frac{A}{H} = \frac{\sqrt{3}}{2}$
- 3 a 4.5 cm                    b 5 cm                    c 3.5 cm
- 4 The ratio to use is the  $\cos \theta = \frac{A}{H}$ .  $\cos 60^\circ = \frac{3.5}{AC}$ . Therefore  $AC = 7$  cm. Iain is correct.
- 5 i-iv  
 In each of the diagrams the smaller the side at the bottom the more each of  $\sin 90^\circ$  tends towards 1,  $\cos 0^\circ$  tends towards 1,  $\sin 0^\circ$  tends towards 0, and  $\cos 90^\circ$  tends towards 0.



- a From the diagram above,  $\tan x = \frac{O}{A} = \frac{1}{1} = 1$ . As the triangle is isosceles each acute angle is equal and therefore  $x = 45$  degrees.
- b The hypotenuse is  $\sqrt{2}$  due to Pythagoras' theorem. Therefore  $\cos x = \frac{A}{H} = \frac{1}{\sqrt{2}}$  and  $\sin x = \frac{O}{H} = \frac{1}{\sqrt{2}}$

**Exercise 22L**

- 1  $65^\circ$
- 2 The safe limits are between 1.04 m and 2.05 m. The highest the ladder will reach is 5.9 m up the wall.
- 3  $44^\circ$
- 4 6.82 m
- 5  $31^\circ$
- 6 a  $25^\circ$   
b 2.10 m  
c Thickness of wood has been ignored
- 7 a  $20^\circ$       b 4.78 m
- 8 She would calculate  $100 \tan 23^\circ$ . The answer is about 42.4 m
- 9 21.1 m
- 10  $\frac{\text{Height}}{3000} = \sin 5.1$ , so height =  $3000 \sin 5.1 = 266.68 = 267$  (to 3 sf)
- 11 8.2 m (2 sf)

**Exercise 22M**

- 1 10.1 km
- 2  $22^\circ$
- 3 429 m
- 4 a 156 m  
b No, the new angle of depression is  $\tan^{-1} \left( \frac{200}{312} \right) = 33^\circ$  and half of  $52^\circ$  is  $26^\circ$
- 5 a 222 m      b  $42^\circ$
- 6 a 21.5 m      b 17.8 m
- 7 13.4 m. We don't know if the angle of elevation is from Sunil's feet or head. This would make a difference to the answer as we would need to add Sunil's height if the angle was from his head.
- 8  $\sin \theta = \frac{1}{3}$  so  $\sin^{-1} 0.3333 = 19.5^\circ$  (3sf)
- 9 The angle is  $16^\circ$  so Cara is not quite correct.

**Exercise 22N**

- 1 a 73.4 km      b 15.6 km
- 2 a 14.7 miles      b 8.5 miles

- 3 Draw a diagram representing the relative places. Your diagram will show the angle of the bearing to fly is  $\tan^{-1} \left( \frac{70}{120} \right) + 90^\circ = 120.256^\circ$  which rounds to  $120^\circ$  (3 sf).
- 4 a 59.4 km      b 8.4 km
- 5 a 15.9 km      b 24.1 km  
c 31.2 km      d  $052^\circ$
- 6 2.28 km
- 7  $235^\circ$
- 8 The distance North is  $120 \cos 35 = 98.298245$  km, so he hasn't gone as far as 100 km, but rounded to the nearest ten he is correct.

**Exercise 22O**

- 1 a 5.79 cm      b  $48.2^\circ$   
c 7.42 cm      d 81.6 cm
- 2 9.86 m
- 3 a  $36.4 \text{ cm}^2$       b  $115 \text{ cm}^2$   
c  $90.6 \text{ cm}^2$       d  $160 \text{ cm}^2$
- 4  $473 \text{ cm}^2$

**Review questions**

- 1 8.6 cm
- 2 6.7 cm
- 3 5.7 cm (1 dp)
- 4 Height =  $5 \sin 75 = 4.8296$  m, which is just higher than 4.8 m so Michael is correct.
- 5  $13.6 \text{ cm}^2$
- 6 2pm
- 7  $237^\circ$
- 8  $52.3^\circ$
- 9 a 5.9 m      b  $82^\circ$
- 10 a  $75.5^\circ$       b  $47.4 \text{ cm}^2$
- 11 The areas are parallelogram  $85.8 \text{ cm}^2$ , triangle  $82.8 \text{ cm}^2$  and circle  $84.9 \text{ cm}^2$ . So all answers are close, but Joe is correct, the parallelogram does have the greatest area.
- 12 a 32.60 m      b 10.07 m  
c 4.10 m      d 9.34 m

## Chapter 23 – Geometry and measures: Congruency and similarity

### Exercise 23A

- 1 a SAS      b SSS      c RHS  
d SSS      e ASA      f ASA
- 2 a i SSS      ii A to R, B to P, C to Q  
b i SAS      ii A to R, B to Q, C to P
- 3 a  $60^\circ$       b  $80^\circ$       c  $40^\circ$       d 5 cm
- 4 The triangle is made of parallel lines so each triangle has a set of corresponding angles with each other. Because the squares share sides with the triangles, each triangle has the same-sized sides. Hence each triangle is congruent, ASA.
- 5 SSS or RHS
- 6 SSS or SAS or RHS
- 7 a always  
b always  
c always  
d sometimes; it's true if the scale factor is 1
- 8 a The square has rotated through angle  $x$ , so each side has rotated the same angle.  
b i ED and DG are both sides of the same square ii CD and AD are both sides of the same square  
c SAS, where CD corresponds to AD, DG corresponds to DE and the angle between them is  $x$  in each case.
- 9 AB and PQ are the corresponding sides to the  $42^\circ$  angle, but they are not equal in length.

### Exercise 23B

- 1 c Each triangle has the same angles as the other.  
d They are all twice as large.  
e 2
- 2 c Each triangle has the same angles as the other  
d They are all three times as large.  
e 3

- 3 c Each triangle has the same angles as the other.  
d yes  
e 1.5
- 4 d Each triangle has the same angles as the other.  
e 2
- 5 d Each triangle has the same angles as the other.  
e 3

### Exercise 23C


- 1 a similar, scale factor 3  
b similar, scale factor 2  
c similar, scale factor 1.5  
d not similar as two corresponding sides have a scale factor of 5, the other one doesn't
- 2 a 3      b angle R      c BA
- 3 a angle B =  $180 - (75 + 50) = 55$  and angle P =  $180 - (55 + 50) = 75$ , so both triangles have the same angles of 50, 55 and 75  
b PR      c BC      d i 28 cm ii 6 cm
- 4 a  $40^\circ$   
b 1.5  
c Multiply length AB by 1.5
- 5 a 8 cm  
b 7.5 cm
- 6 a i ABC with DEF  
ii 3  
iii  $x = 3$  cm  $y = 18$  cm  
b i GHI with LKJ  
ii 0.5  
iii  $x = 8$  cm  $y = 3.5$  cm  
c i PQR with TUS  
ii 1.5  
iii  $x = 5.33$  cm  $y = 12$  cm  
d i MNL with WUV  
ii 5  
iii  $x = 9$  cm  $y = 40$  cm
- 7 a Sides in same ratio  
b 3      c 13 cm      d 39 cm

- 8 a The interior angles of all regular hexagons are  $120^\circ$ , so all the angles of each hexagon will correspond, hence similar shapes.  
 b 12 cm  
 c 6.928 cm

### Exercise 23D

- 1 82 m  
 2  $\frac{\text{pole}}{330} = \frac{400}{600}$ , pole =  $330 \times \frac{4}{6} = 220$  feet  
 3 15 m  
 4 3.3 m


- 5 a



- b 1.8 m

- 6 a i ABC and ADE                    ii 9 cm  
 b i ABC and ADE                    ii 12 cm

- 7 a



- b  $\frac{CD}{9} = \frac{9}{6}$ ,  $CD = 9 \times \frac{9}{6} = 13.5$  cm

- 8 c 4 cm

- iv enlargement  
 v no transformation  
 c It is not true because E, the enlarged triangle, is not congruent, and nor is F.

- 2 Ellie is correct:  
 a both triangles have an identical set of angles: 90, 65 and 25  
 b both shapes are rectangles and the lengths of the large rectangle are four times larger than the small one, so the large rectangle is an enlargement of the small one.

- 3 a always  
 b always  
 c sometimes  
 d sometimes  
 e always

- 4 a congruent SAS  
 b congruent RHS  
 c not congruent  
 d congruent SSS

- 5 Scale factor between similar triangles is  $1000 \div 40 = 25$ , hence tree height =  $140 \times 25 = 3500$  cm = 35 m

- 6 Construction of a triangle with sides exactly the same size as the given triangle

- 7 ABX with DCX and ADX with BCX

- 8 Andrew is correct. Eve has calculated the length of AD (8 cm) so ED should be 2 cm.

- 9  $4.5 \text{ cm}^2$

### Review questions

- 1 a B, C and D  
 b i rotation  
 ii reflection  
 iii reflection

## Chapter 24 – Probability: Combined events

### Exercise 24A

- 1 a 7    b 2 and 12  
c

Score	Probability
2	$\frac{1}{36}$
3	$\frac{1}{18}$
4	$\frac{1}{12}$
5	$\frac{1}{9}$
6	$\frac{5}{36}$
7	$\frac{1}{6}$
8	$\frac{5}{36}$
9	$\frac{1}{9}$
10	$\frac{1}{12}$
11	$\frac{1}{18}$
12	$\frac{1}{36}$

- d i  $\frac{1}{12}$       ii  $\frac{1}{3}$                           iii  $\frac{1}{2}$   
iv  $\frac{7}{36}$       v  $\frac{5}{12}$                                   vi  $\frac{5}{18}$

2 a  $\frac{1}{12}$       b  $\frac{11}{36}$       c  $\frac{1}{6}$     d  $\frac{5}{9}$

3 a  $\frac{1}{36}$       b  $\frac{11}{36}$       c  $\frac{5}{18}$

4 a  $\frac{1}{4}$       b  $\frac{1}{2}$       c  $\frac{3}{4}$     d  $\frac{1}{4}$

5 a  $\frac{1}{12}$       b  $\frac{1}{4}$       c  $\frac{1}{6}$

6 a

6	5	4	3	2	1	0
5	4	3	2	1	0	1
4	3	2	1	0	1	2
3	2	1	0	1	2	3
2	1	0	1	2	3	4
1	0	1	2	3	4	5
	1	2	3	4	5	6

Score on second dice

Score on first dice

- b i  $\frac{5}{18}$     ii  $\frac{1}{6}$     iii  $\frac{1}{9}$     iv 0    v  $\frac{1}{2}$

7 a

5	6	7	8	9	10
4	5	6	7	8	9
3	4	5	6	7	8
2	3	4	5	6	7
1	2	3	4	5	6
	1	2	3	4	5

Score on second spinner

Score on first spinner

- b 6  
c i  $\frac{4}{25}$     ii  $\frac{13}{25}$     iii  $\frac{1}{5}$     iv  $\frac{3}{5}$

8  $\frac{18}{64} = \frac{1}{8}$

9 It will show all the possible products.

10 a HHH, HHT, HTH, THH, HTT, THT, TTH, TTT

- b i  $\frac{1}{8}$     ii  $\frac{3}{8}$     iii  $\frac{1}{8}$     iv  $\frac{7}{8}$

11 a 16      b 32      c 1024      d  $2^n$

12 a

	1	2	3	4	5	6
1	2	3	4	5	6	7
2	3	4	5	6	7	8
3	4	5	6	7	8	9
4	5	6	7	8	9	10
5	6	7	8	9	10	11
6	7	8	9	10	11	12

- b  $\frac{1}{18}$     c 18    d twice



## Exercise 24B

1 a

		Shaded	Unshaded
Shape	Circle	3	3
	Triangle	2	2

b i  $\frac{1}{5}$                       ii  $\frac{1}{2}$ 2 a 40                      b 45%                      c  $\frac{3}{40}$ 

3 a

	England	Scotland	Wales	Total
Female	26	12	4	42
Male	24	8	6	38
Total	50	20	10	80

b  $\frac{1}{4}$                                       c  $\frac{3}{19}$ 

4 a

	Blackpool	Chester Zoo	Total
Boys	32	28	60
Girls	26	14	40
Total	58	42	100

b  $\frac{29}{50}$                                       c  $\frac{7}{20}$ 5 a 23                      b 20%                      c  $\frac{3}{50}$                       d 520b 9                      c 0                      d  $\frac{5}{12}$ 

e The 0 on the first spinner cannot be taken as odd.

6 a 15%                      b  $\frac{9}{40}$                       c 75

7 a

Type of rose	Colour of rose					
	White	Red	Orange	Yellow	Pink	Copper
Dwarf						
Small						
Medium						
Large climbing						
Rambling						


b i  $\frac{1}{5}$                                       ii  $\frac{4}{5}$ 8 a  $\frac{2}{45}$                       b 40%                      c 45%

d No, as you don't know how much the people who get over £350 actually earn


## Exercise 24C

1 a 0.9                                      b 0.7


2 a 0.75                                      b 0.45

b i  $\frac{2}{5}$     ii  $\frac{3}{5}$     iii  $\frac{1}{2}$     iv  $\frac{1}{2}$     v  $\frac{7}{10}$     vi  $\frac{1}{5}$ 4 a i 0.52                      ii 0.48                      iii 0.65                      iv 0.35  
v 0.82                      vi 0.35

b 0.3



b i 0.5                      ii 0.6                      iii 0.3




b i 0.65                      ii 0.7                      iii 0.9                      iv 0.45

7 a 130

b i  $\frac{8}{13}$ 

ii The probability that a student chosen at random walks to and from school


c  $\frac{5}{26}$ a  $\frac{1}{2}$                       b  $\frac{2}{5}$                       c  $\frac{7}{10}$                       d  $\frac{1}{5}$ 

9 0.4

10 0.5



11 a  $(A \cup B)'$                       b  $(A \cap B)'$

12




13  $\frac{37}{80}$

**Exercise 24D**




2 a i  $\frac{1}{4}$     ii  $\frac{1}{2}$     iii  $\frac{3}{4}$

3 a  $\frac{1}{13}$     b  $\frac{12}{13}$



d i  $\frac{1}{169}$     ii  $\frac{24}{169}$     iii  $\frac{25}{169}$


4 a  $\frac{2}{3}$     b  $\frac{1}{2}$




d i  $\frac{1}{6}$     ii  $\frac{1}{2}$     iii  $\frac{5}{6}$

e 15 days


5 a  $\frac{2}{5}$



i  $\frac{4}{25}$     ii  $\frac{12}{25}$



- b i**  $\frac{1}{8}$       **ii**  $\frac{3}{8}$       **iii**  $\frac{7}{8}$



- b** 0.14      **c** 0.41      **d** 0.09

- 8 a** 1      **b** 1      **c**
- |  | First event | Second event | Outcome | Probability                      |
|--|-------------|--------------|---------|----------------------------------|
|  | A           | C            | (A, C)  | $\frac{2}{5} \times \frac{3}{4}$ |
|  | A           | D            | (A, D)  | $\frac{2}{5} \times \frac{1}{4}$ |
|  | B           | E            | (B, E)  | $\frac{3}{5} \times \frac{1}{3}$ |
|  | B           | F            | (B, F)  | $\frac{3}{5} \times \frac{2}{3}$ |

**b**

$$= \frac{3}{10}$$

$$= \frac{1}{10}$$

$$= \frac{1}{5}$$

$$= \frac{2}{5}$$

- 9 a** 27      **b** 3
- 10 a**  $\frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{16}$       **b**  $\frac{1}{2}n$

## Review questions

- 1 a**


1		Spinner				
		2	3	4	5	
Coin	H	1, H	2, H	3, H	4, H	5, H
	T	1, T	2, T	3, T	4, T	5, T

- b**  $\frac{3}{10}$

● ANSWERS


2 a 23      b 20%      c  $\frac{4}{25}$       d 480

3 a 110      b  $\frac{19}{55}$       c  $\frac{3}{22}$       d  $\frac{36}{55}$




5 a 0.1

b i 0.3      ii 0.5      iii 0.6      iv 0.3




b i  $\frac{70}{260} = 0.269$       ii  $\frac{60}{260} = 0.231$

iii  $\frac{5}{260} = 0.0192$       iv  $\frac{20}{165} = 0.121$



b  $\frac{5}{9}$

c  $\frac{4}{9}$



b i 0.88      ii 0.12

c calculate  $0.88^2$

## Chapter 25 – Number: Powers and standard form

### Exercise 25A

1 a  $2^4$       b  $3^5$       c  $7^2$       d  $5^3$   
 e  $10^7$       f  $6^4$       g  $4^1$       h  $1^7$   
 i  $0.5^4$       j  $100^3$

2 a  $3 \times 3 \times 3 \times 3$   
 b  $9 \times 9 \times 9$   
 c  $6 \times 6$   
 d  $10 \times 10 \times 10 \times 10 \times 10$   
 e  $2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2$   
 f 8  
 g  $0.1 \times 0.1 \times 0.1$   
 h  $2.5 \times 2.5$   
 i  $0.7 \times 0.7 \times 0.7$   
 j  $1000 \times 1000$

3 a 16      b 243  
 c 49      d 125  
 e 10 000 000      f 1296  
 g 4      h 1  
 i 0.0625      j 1 000 000

4 a 81      b 729  
 c 36      d 100 000  
 e 1024      f 8  
 g 0.001      h 6.25  
 i 0.343      j 1 000 000

5  $125 \text{ m}^3$

6 b  $10^2$       c  $2^3$       d  $5^2$

7 a 1      b 4      c 1  
 d 1      e 1

8 Any power of 1 is equal to 1.

9  $10^6$

10  $10^6$

11 a 1      b -1      c 1  
 d 1      e -1

12 a 1      b -1      c -1  
 d 1      e 1

13  $2^{24}, 4^{12}, 8^8, 16^6$

14  $x = 7$

### Exercise 25B

1 a  $5^4$       b  $5^{10}$       c  $5^5$       d  $5^3$   
 e  $5^{15}$       f  $5^9$       g  $5^6$       h  $5^9$   
 i  $5^8$

2 a  $x^8$       b  $x^9$       c  $x^8$       d  $x^5$   
 e  $x^{12}$       f  $x^{13}$       g  $x^{11}$       h  $x^{10}$   
 i  $x^{16}$

3 a  $6^3$       b  $6^5$       c  $6^1$       d  $6^0$   
 e  $6^1$       f  $6^3$       g  $6^2$       h  $6^1$   
 i  $6^2$

4 a  $x^4$       b  $x^5$       c  $x^3$       d  $x^3$   
 e  $x^6$       f  $x^5$       g  $x^2$       h  $x^6$   
 i  $x^9$

5 a 1      b  $6^0$       c 1

6 a 1      b  $5^0$       c 1

7 Answers for power 0 are always 1.

8 Two values with a sum of 7, for example  $a = 5$  and  $b = 2$

9 a  $a = 1$  and  $x = 2$   
 b  $x = 1$

10 a  $6a^5$       b  $9a^2$       c  $8a^6$   
 d  $-6a^4$       e  $8a^8$       f  $-10a^{-3}$

11 a  $3a$       b  $4a^3$       c  $3a^4$   
 d  $6a$       e  $4a^3$       f  $5a^{-4}$

12 a Possible answer:  $6x^2 \times 2y^5$  and  $3xy \times 4xy^4$   
 b Possible answer:  $24x^2y^7 \div 2y^2$  and  $12x^6y^8 \div x^4y^3$

13 a  $3a^3b^2$       b  $3a^2c^4$       c  $8a^2b^2c^3$

14 a  $8a^5b^4$       b  $10a^3b$       c  $30a^{-2}b^{-2}$   
 d  $2ab^3$       e  $8a^{-5}b^7$       f  $4a^5b^{-5}$

15 a  $4^6$       b  $4^{15}$       c  $4^6$   
 d  $4^{-6}$       e  $4^6$       f  $4^0$

16  $12$  ( $a = 2, b = 1, c = 3$ )

### Exercise 25C

1 a 60 000      b 120 000      c 150  
 d 42 000      e 1400      f 300  
 g 400      h 8000      i 160 000  
 j 4500      k 8000      l 250 000

2 a 5      b 50      c 25      d 30  
 e 7      f 300      g 6      h 30  
 i 1      j 15      k 40      l 5  
 m 40      n 320

3 a 54 400      b 16 000

4  $30 \times 90\,000 = 2\,700\,000$   
 $600 \times 8000 = 4\,800\,000$   
 $5000 \times 4000 = 20\,000\,000$   
 $200\,000 \times 700 = 140\,000\,000$

- 5 1400 million
- 6 a 31 b 310 c 3100 d 31 000
- 7 a 65 b 650 c 6500 d 65 000
- 8 a 0.31 b 0.031 c 0.0031 d 0.000 31
- 9 a 0.65 b 0.065 c 0.0065 d 0.000 65
- 10 a 250 b 34.5 c 4670  
d 346 e 207.89 f 56 780  
g 246 h 0.76 i 999 000  
j 23 456 k 98 765.4 l 43 230 000  
m 345.78 n 6000 o 56.7  
p 560 045
- 11 a 0.025 b 0.345  
c 0.004 67 d 3.46  
e 0.207 89 f 0.056 78  
g 0.0246 h 0.0076  
i 0.000 000 999 j 2.3456  
k 0.098 765 4 l 0.000 043 23  
m 0.000 000 034 578 n 0.000 000 000 06  
o 0.000 000 567 p 0.005 600 45
- 12 a 230 b 578 900 c 4790  
d 57 000 000 e 216 f 10 500  
g 0.000 32 h 9870

13 They are all true.

### Exercise 25D

- 1 a 0.31 b 0.031 c 0.0031 d 0.000 31
- 2 a 0.65 b 0.065 c 0.0065 d 0.000 65
- 3 a  $9999999999 \times 10^{99}$   
b  $0.000000001 \times 10^{-99}$  (depending on number of digits displayed)
- 4 a 31 b 310 c 3100 d 31 000
- 5 a 65 b 650 c 6500 d 65 000
- 6 a 250 b 34.5 c 0.004 67  
d 34.6 e 0.020 789 f 5678  
g 246 h 7600 i 897 000  
j 0.008 65 k 60 000 000 l 0.000 567
- 7 a  $2.5 \times 10^2$  b  $3.45 \times 10^{-1}$   
c  $4.67 \times 10^4$  d  $3.4 \times 10^9$   
e  $2.078 \times 10^{10}$  f  $5.678 \times 10^{-4}$   
g  $2.46 \times 10^3$  h  $7.6 \times 10^{-2}$   
i  $7.63 \times 10^{-4}$  j  $9.99 \times 10^{-1}$   
k  $2.3456 \times 10^2$  l  $9.87654 \times 10$   
m  $6 \times 10^{-4}$  n  $5.67 \times 10^{-3}$   
o  $5.60045 \times 10$

- 8  $2.7797 \times 10^4$
- 9  $2.81581 \times 10^5$ ,  $3 \times 10$ ,  $1.382101 \times 10^6$
- 10  $1.298 \times 10^7$ ,  $2.997 \times 10^9$ ,  $7.4 \times 10^4$
- 11 a  $5.67 \times 10^3$  b  $6 \times 10^2$  c  $3.46 \times 10^{-1}$   
d  $7 \times 10^{-4}$  e  $5.6 \times 10^2$  f  $6 \times 10^5$   
g  $7 \times 10^3$  h 1.6 i  $2.3 \times 10^7$   
j  $3 \times 10^{-6}$  k  $2.56 \times 10^6$  l  $4.8 \times 10^2$   
m  $1.12 \times 10^2$  n  $6 \times 10^{-1}$  o  $2.8 \times 10^6$
- 12 a  $1.08 \times 10^8$  b  $4.8 \times 10^6$  c  $1.2 \times 10^9$   
d 1.08 e  $6.4 \times 10^2$  f  $1.2 \times 10$   
g 2.88 h  $2.5 \times 10^7$  i  $8 \times 10^{-6}$
- 13  $2 \times 10^{13}$ ,  $1 \times 10^{-10}$ , mass =  $2 \times 10^3$  g (2 kg)
- 14  $5 \times 10^4$
- 15 7800,  $3.7 \times 2.1 \times 10^4$ ,  $7.85 \times 10^4$ ,  $79 \times 10^3$
- 16 0.0005,  $4 \times 10^{-3}$ ,  $0.52 \times 10^{-2}$ ,  $6 \times 10^{-2}$
- 17 a  $1.8 \times 10^4$  b  $1.7 \times 10^7$  c  $8.8 \times 10^7$   
d  $1.9 \times 10^5$  e  $6.9 \times 10^6$  f  $7.1 \times 10^{-9}$   
g  $1.2 \times 10^{-7}$  h  $5.3 \times 10^8$
- 18 a  $1 \times 10^0$   
b  $1.0 \times 10$   
c It is easier to use ordinary numbers in this case

### Review questions

- 1 a 64 b 64 c 10 000 000
- 2 a 62 b 4900 c 3.47 d 0.123
- 3 a 6 000 b 300 000 c 80 d 40
- 4 a 196 b units digits is  $5 \times 5 = 25$  so it should end in 5
- 5 a  $7^9$   
b  $x^4$   
c He multiplied the powers of x, but he should have added them.  $15x^9$
- 6 a  $t^8$  b  $m^5$  c  $9x^6$  d  $10a^7h^5$
- 7 a  $x^{11}$  b  $m^{-5}$  c  $8k^5m^3$
- 8 a  $7.5 \times 10^4$  b 0.009
- 9 a 2y b  $8m^9p^{12}$
- 10 a 1200  
b  $1 \times 10^b = 1^b \times 10^b$  or  $a \times 10^1 = a^1 \times 10^1$
- 11 3

## Chapter 26 – Algebra: Simultaneous equations and linear inequalities

### Exercise 26A

- 1 a  $x = 4\frac{1}{2}, y = 1\frac{1}{2}$       b  $x = -2, y = 4$   
 c  $x = 2\frac{1}{2}, y = -1\frac{1}{2}$       d  $x = 3, y = 10$   
 e  $x = 9, y = -2$       f  $x = 0.5, y = 4$
- 2 a  $a = 7, b = 10$       b  $c = 4, d = 11$   
 c  $e = 5, f = 3$       d  $g = -1, h = 6$   
 e  $j = 13, k = 2$       f  $m = 1.5, n = -7$
- 3  $x = 12, y = 2$

### Exercise 26B

- 1 a  $x = 9, y = -2$       b  $x = \frac{1}{2}, y = 5$   
 c  $x = -3, y = -10$       d  $x = -12, y = -6$   
 e  $x = 3, y = 5$       f  $x = 0.5, y = 6.5$
- 2 a  $x = 2\frac{1}{4}, y = 6\frac{1}{2}$       b  $x = 4, y = 3$   
 c  $x = 5, y = 3$       d  $x = 4, y = 5$   
 e  $x = 11, y = -3$       f  $x = -7, y = 2$
- 3 a  $x = 1, y = 3$       b  $x = 5, y = 9$

### Exercise 26C

- 1 a  $x = 2, y = 5$       b  $x = 4, y = -3$   
 c  $x = 1, y = 7$       d  $x = \frac{1}{2}, y = -\frac{3}{4}$   
 e  $x = -1, y = 5$       f  $x = 1\frac{1}{2}, y = \frac{3}{4}$
- 2 a  $x = 5, y = 1$       b  $x = 3, y = 8$   
 c  $x = 9, y = 1$       d  $x = 7, y = 3$   
 e  $x = 4, y = 2$       f  $x = 6, y = 5$
- 3 a  $a = 10, b = \frac{1}{2},$       b  $c = -3, d = 5,$   
 c  $e = 1, f = 14,$       d  $g = -2, h = -5,$   
 e  $m = 8, n = 0,$       f  $p = 3\frac{1}{2}, q = -2,$   
 g  $r = 7, s = 7,$       h  $t = \frac{2}{3}, u = -1,$   
 i  $v = 56, w = 32$

### Exercise 26D

- 1 a  $14a + b = 47$       b  $a = 3, b = 5$   
 c 146, 443
- 2 a  $3t + 5b = 810, 3t + 3b = 630$   
 b i £1.20      ii 90p  
 c £10.20
- 3 a  $10x + 5y = 840, 8x + 10y = 1044$   
 b £4.07

- 4  $(1, -2)$  is the solution to equations A and C;  
 $(-1, 3)$  is the solution to equations A and D;  
 $(2, 1)$  is the solution to B and C;  $(3, -3)$  is  
 the solution to B and D.

- 5 84p
- 6 10.3 kg
- 7  $p = 36, c = 22$ . Total weight for Baz is 428  
 pounds so he can carry the load safely on  
 his trailer.
- 8 £4.40
- 9  $b = £3.50, p = £1.75$ . Camilla needs £35 so  
 she will not have enough money.
- 10 a My age minus 6 equals 2  $x$  (my son's age  
 minus 6)  
 b  $x = 46$  and  $y = 26$
- 11 Amul £7.20, Kim £3.50
- 12 a They are the same equation. Divide the  
 first by 2 and it is the second, so they  
 have an infinite number of solutions.  
 b Double the second equation to get  
 $6x + 2y = 14$  and subtract to get  $9 = 14$ .  
 The left-hand sides are the same if the  
 second is doubled so they cannot have  
 different values.






















### Exercise 26E

- 1 a  $x < 3$       b  $t > 8$       c  $p > 10$       d  $x < 5$   
 e  $y < 3$       f  $t > 5$       g  $x < 6$       h  $y < 15$   
 i  $t > 18$       j  $x < 7$       k  $x < 3$       l  $t > 5$
- 2 a 7      b 6      c 16  
 d 3      e 7
- 3 a 11      b 16      c 16  
 d 3      e 7
- 4  $2x + 3 < 20, x < 8.50$ , so the most each could  
 cost is £8.49
- 5 a Because  $3 + 4 = 7$ , which is less than the  
 third side of length 8  
 b  $x + x + 2 > 10, 2x + 2 > 10, 2x > 8, x > 4$ , so  
 smallest value of  $x$  is 5
- 6 a  $x = 6$  and  $x < 3$  scores  $-1$  (nothing in  
 common),  $x < 3$  and  $x > 0$  scores  $+1$  (1 in  
 common for example),  $x > 0$  and  $x = 2$   
 scores  $+1$  (2 in common),  $x = 2$  and  $x > 4$   
 scores  $-1$  (nothing in common), so we  
 get  $-1 + 1 + 1 - 1 = 0$





- b**  $x > 0$  and  $x = 6$  scores +1 (6 in common),  
 $x = 6$  and  $x > 4$  scores +1 (6 in common),  
 $x > 4$  and  $x = 2$  scores -1 (nothing in  
common),  $x = 2$  and  $x < 3$  scores +1 (2 in  
common).  $+1 + 1 - 1 + 1 = 2$
- c** Any acceptable combination, e.g.  $x = 2$ ,  
 $x < 3, x > 0, x > 4, x = 6$

- 7** **a**  $y \leq 4$     **b**  $x \geq -2$     **c**  $x \leq \frac{14}{5}$   
**d**  $x > 38$     **e**  $x < 6\frac{1}{2}$     **f**  $y \leq \frac{7}{5}$
- 8** **a**  $3 < x < 6$     **b**  $2 < x < 5$     **c**  $-1 < x \leq 3$   
**d**  $1 \leq x < 4$     **e**  $2 \leq x < 4$     **f**  $0 \leq x \leq 5$
- 9** 6 bottles

### Exercise 26F







- 1** **a**  $x > 1$     **b**  $x \leq 3$     **c**  $x < 2$   
**d**  $x \geq -1$     **e**  $1 < x \leq 4$     **f**  $-2 < x < 4$
- 2** **a**     **b**   
**c**     **d**   
**e**     **f**   
**g**     **h** 
- 3** **a**  $x \geq 4$      **b**  $x < -2$    
**c**  $x \leq 3$      **d**  $x > 3$    
**e**  $x > 4$      **f**  $x \leq 5$    
**g**  $x < 1.5$      **h**  $x \leq -2$    
**i**  $x > 50$      **j**  $x \leq 2.5$    
**k**  $x < 10$      **l**  $x > 13$  
- 4** **a**  $x > 22$   
**b**  $x \leq 25$   
**c** 
- d** Apples could cost 23p, 24p or 25p.


- 5** 1 and 4

- 6** **a**  $x > 2$    
**b**  $x \geq 6$    
**c**  $x \leq -1$    
**d**  $x \geq -4$  

### Review questions

- 1** **a**  $x > 23$ ,    **b**  $x \leq 15$ ,    **c**  $x < 48$ ,  
**d**  $x \geq 6$ ,    **e**  $5 \leq x \leq 25$

- 2** **a**  $x < 7$      **b**  $x > 15$    
**c**  $x \leq 2$      **d**  $x \geq 4$    
**e**  $x \geq -12$      **f**  $x < 8$  


- 3** 

- 4** 9

- 5** £340

- 6** 2 hr 10 min

- 7** **i**  $-3 < x < 1$ , number line B;    **ii**  $-2 < x < 4$ ,  
number line below;    **iii**  $-1 < x < 2$ , number  
line A



- 8**  $x = 36, y = 216$

- 9** £195

- 10** **a**  $x = 4, y = 3$   
**b** **i**  $1000x + 1000y = 7000 \rightarrow x + y = 7$   
**ii**  $984x - 984y = 984 \rightarrow x - y = 1$   
**c**  $a = 9, b = 5$

- 11**  $x = 5, y = -1$



- 12 Let straight part of track =  $D$ , inner radius of end =  $r$ , outer radius =  $r + x$   
 $x$  being the width of the track  
 Length of inner track =  $2D + 2\pi r = 300$   
 ..... i  
 Length of outer track =  $2D + 2\pi(r + x) = 320$   
 .....ii  
 Subtract equation i from ii to give

$$\begin{aligned} 2\pi(r + x) - 2\pi r &= 20 \\ 2\pi r + 2\pi x - 2\pi r &= 20 \\ 2\pi x &= 20 \\ x &= 3.2 \text{ 2sf} \end{aligned}$$

## Chapter 27 – Algebra: Non-linear graphs

### Exercise 27A

- 1 a 1 hour                      b  $1\frac{1}{2}$  hours  
 c  $1\frac{1}{2}$  hours                d 45 km per hour  
 e 30 km per hour
- 2 a i 2 h            ii 3 h            iii 5 h  
 b i 40 km/h    ii 120 km/h    iii 40 km/h
- 3 a  $2\frac{1}{2}$  km/h    b 3.75 m/s    c  $2\frac{1}{2}$  km/h
- 4 a 30 km        b 40 km        c 50 km  
 d 100 km/h
- 5 a i 263 m/min (3 sf)    ii 15.8 km/h (3 sf)  
 b i 500 m/min  
 c Paul by 1 minute
- 6 a Patrick ran quickly at first, then had a slow middle section but he won the race with a final sprint. Araf ran steadily all the way and came second. Sean set off the slowest, speeded up towards the end but still came third.  
 b i 1.67 m/s    ii 6 km/h
- 7 There are three methods for doing this question. This table shows the first, which is writing down the distances covered each hour.

Time	Walker	Cyclist
9am	0	0
9:30	3	0
10:00	6	0
10:30	9	0
11:00	12	7.5
11:30	15	15
12:00	18	22.5
12:30	21	30


The second method is algebra:

Walker takes  $T$  hours until overtaken, so  $T = \frac{D}{6}$ ;

Cyclist takes  $T - 1.5$  to overtake, so  $T - 1.5 = \frac{D}{15}$ .

Rearranging gives  $15T - 22.5 = 6T$ ,  $9T = 22.5$ ,  $T = 2.5$ .

The third method is a graph:



All methods give the same answer of 11:30 when the cyclist overtakes the walker.

- 8 a Vehicle 2 overtook Vehicle 1  
 b Vehicle 1 overtook Vehicle 2  
 c Vehicles passed travelling in different directions  
 d Vehicle 2 overtook Vehicle 1  
 e 17:15  
 f 11.3 mph if you only count travelling time, or 5.1 mph if you count total time.

### Exercise 27B

- 1 a Two taps on  
 b One tap on  
 c Maureen gets in the bath  
 d Maureen has a bath  
 e Maureen takes the plug out, water leaves the bath  
 f Maureen gets out of the bath  
 g Water continues to leave the bath until the bath is empty
- 2 a Graph C
- b A B D
- 3 a b c   
 d e f   
 g

### Exercise 27C

1

$x$	-5	-4.5	-4	-3.5	-3	-2.5
$y$	25	20.25	16	12.25	9	6.25
		-2	-1.5	-1	-0.5	0
		4	2.25	1	0.15	0

2

$x$	-3	-2	-1	0	1	2	3
$y$	27	12	3	0	3	12	27

3 a

$x$	-5	-4	-3	-2	-1	0	1	2	3	4	5
$y$	27	18	11	6	3	2	3	6	11	18	27

b  $y = 2$

4 a

$x$	-5	-4	-3	-2	-1	0	1	2	3	4	5
$x^2$	25	16	9	4	1	0	1	4	9	16	25
$-3x$	15	12	9	6	3	0	-3	-6	-9	-12	-15
$y$	40	28	18	10	4	0	-2	-2	0	4	10

b 1.8      c -1.2, 4.2      d  $y = 0$

5 a Values of  $y$ : 7, 0, -5, -8, -9, -8, -5, 0, 7


b -6.7

c -0.45 or 4.45

6 a Values of  $y$ : 15, 8, 3, 0, -1, 0, 3, 8, 15

b -0.75      c -1.6, -4.4

d



e (-4, 0) and (-1.5, 1.25)

7 a  $x$ : -2, -1, 0, 1, 2, 3, 4

$y$ : 8, 3, 0, -1, 0, 3, 8

b (0, 0) and (3, 3)

8 a They have the same value,  $x = 4$

b

$x$	-3	-2	-1	0	1	2	3	4	5	6	7
$y$	28	19	12	7	4	3	4	7	12	19	28

Since the quadratic graph has a vertical line of symmetry and the  $y$ -values for  $x = 1$  and  $x = 3$  are the same, this means that the  $y$ -values will be symmetric about  $x = 2$ . Hence the  $y$ -values will be the same for  $x = 0$  and  $x = 4$ , and so on.

9 a  $x$ : -4, -3, -2, -1, 0, 1, 2, 3, 4

$y$ : 12, 5, 0, -3, -4, -3, 0, 5, 12


b (2, 0) and (-2, 0)

c -1.75

d  $x = 3.5$  or  $-3.5$

10 a

$x$	-4	-3	-2	-1	0	1	2	3	4
$y$	-12	-5	0	3	4	3	0	-5	-12



b (2, 0) and (-2, 0)

c 1.75

d  $x = 3.5$  or  $-3.5$

e The curves are reflections in the  $x$ -axis.




## Exercise 27D

- 1 a  $-2, -5$  b  $4, 9$  c  $-6, 3$   
 d  $-3, -1$  e  $-6, -4$  f  $-3, 2$   
 g  $-1, 3$  h  $-4, 5$  i  $1, -2$   
 j  $2, -5$  k  $7, -4$  l  $3, 2$   
 m  $1, 5$  n  $4, 3$  o  $-7, 5$
- 2 a  $-2, -3$  b  $-4, -1$  c  $2, 4$   
 d  $-2, 5$  e  $-3, 5$  d:  $-3, 5$  f  $-6, 3$   
 g  $-1, 2$  h  $-2, -5$  i  $7, 7$   
 j  $-9, -2$  k  $3, 5$  l  $-6, 2$   
 m  $-2, -2$  n  $4$  o  $-2, -6$
- 3 a  $x^2 + 6x - 55 = 25 + 30 - 55 = 0$   
 b  $x = -11$
- 4 a  $5, -10$  b  $7, -8$  c  $1, 18$   
 d  $-2, -11$  e  $2, -15$  f  $3, 10$   
 g  $-1, -999$  h  $1$  i  $-10, 1$
- 5  $x^2 + 8x + 16 = (x + 2)(x + 6) = 0$  so the answers are  $-2$  and  $-6$
- 6  $x^2 + 5x - 6 = (x + 6)(x - 1) = 0$  so the answers are  $1$  and  $-6$
- 7 a Area  $= (x + 1)(x + 2) = x^2 + 3x + 2 = 42$ ,  
 therefore  $x^2 + 3x - 40 = 0$   
 b  $x = 5$  cm (discard the negative value as  $x$  is a length)  
 c Perimeter  $= 26$  cm

8  $x = 9$  cm

## Exercise 27E

- 1 a  $(0, -5), (0, 8)$  and  $(0, 0)$ .  
 b Each equation is written in the form  $x^2 + ax + b$ . You should find that the  $y$ -intercept is the value of  $b$ .




- 2 a  $x = -1$  or  $5$ ;  $x = -2$  or  $-4$ ;  $x = 0$  or  $2$   
 b  $(-1, 0), (5, 0); (-4, 0), (-2, 0); (0, 0), (2, 0)$
- 3  $(2, -9), (-3, -1), (1, -1)$


## Exercise 27F

- 1 i  $(0, 5)$  ii  $(-5, 0)$  and  $(1, 0)$   
 iii  $(-2, 9)$
- 2 i  $(0, -3)$  ii  $(-1, 0)$  and  $(3, 0)$   
 iii  $(1, -4)$
- 3 a roots:  $(-2, 0)$  and  $(2, 0)$ ;  $y$ -intercept  $(0, -4)$   
 b roots:  $(0, 0)$  and  $(6, 0)$ ;  $y$ -intercept  $(0, 0)$   
 c roots:  $(-1, 0)$  and  $(3, 0)$ ;  $y$ -intercept  $(0, -3)$   
 d roots:  $(-11, 0)$  and  $(-3, 0)$ ;  $y$ -intercept  $(0, 33)$
- 4  $(2, -1)$
- 5 i  $(0, 9)$  ii  $(3, 0)$  and  $(-3, 0)$  iii  $(0, 9)$
- 6 i  $(0, 16)$  ii  $(2, 0), (8, 0)$  iii  $(5, -9)$
- 7 i  $(0, -15)$   
 ii  $(3, 0), (-5, 0)$   
 iii  $(-1, -16)$
- 8 i  $(0, -6)$   
 ii  $(-2, 0), (1.5, 0)$   
 iii  $(-0.25, -6.1)$

**Exercise 27G**



**2 a** Values of  $y$ :  $-26, -7, 0, 1, 2, 9, 28$




**b** 16.6

**3 a** Values of  $y$ :  $-24, -12.63, -5, -0.38, 2, 2.9, 3, 3.13, 4, 6.38, 11, 18.63, 30$

**b** 4.7      **c**  $(-1.45, 0)$


**4 a**



Values of  $y$ :  $-0.05, -0.1, -0.2, -0.25, -0.5, -1, -2, -2.5, -5, 5, 2.5, 2, 1, 0.5, 0.2, 0.1, 0.067, 0.05$

**b**  $\frac{1}{0}$  is undefined

**c**



d 0.4 and 2.6

- 5 a Quadratic    b Linear    c None  
 d Reciprocal    e None    f Cubic  
 g Linear        h None    i Quadratic

## Review questions

1 a




b Answers will vary

2 a 160 km/h

b 80 km/h

c




d 14:41

3  $x = 9$  or  $10$

4 23, 10, 5, 2

5 a



b i 0.4, 2.6

ii  $-0.3, 3.3$

iii  $-1.4, 4.4$

6 a  $x = 6$     b 38 cm

7

