


Guidance on the use of codes for this mark scheme	
M	Method mark
A	Accuracy mark
B	Accuracy mark in AO1
C	Communication mark
P	Proof or process mark
oe	Or equivalent
ft	Follow through
cao	Correct answer only



Question	Working	Answer	Mark	AO	Notes	Grade
1 a b c d e f		$D = 7w$ $C = pn$ $Y = \frac{m}{12}$ $P = 100D$ $A = lw$ $P = nl$	B1 B1 B1 B1 B1 B1	3	B1 oe B1 oe B1 oe B1 oe B1 oe B1 oe	B
2 a b		$C = 80h$ $C = 80h + 50$	B1 B1	3	B1 cao B1 cao	B
3 a b		<p>No To be able to work out what the number thought of, you need to know the answer.</p> <p>Yes Because I can write an equation from the information and solve it. $x + 15 = 26$ so $x = 11$</p>	C1 C1 B1	2	C1 for No and a reason C1 for Yes and a reason B1 for showing the equation and the solution	B
4		<p>For example, in the rule $\text{pay} = 15 \times$ hours. As hours varies, so will the calculation to calculate pay. Yes, there will be others, there will be hundreds of different possible calculations.</p>	C1 C1 C1	2	C1 for an explanation of why it is possible for more than one calculation to match with the same rule C1 for using an example to go alongside the explanation C1 for stating Yes there will be more, and qualifying this	B
5		<p>We use (x, y) to describe the position, where the first part, x, is along the x-axis. Then the second part, y, is along the y-axis. Example, e.g. The convention for point A is $(2, 3)$. If we didn't have the convention then we could use $(3, 2)$ but that could be confused now with point B.</p>	C1 C1	2	C1 for clear explanation C1 for a clear example illustrated with a sketch graph	B

6	a	Yes For example we could write as $2x = y - 6$ Rearranging an equation.	C1	2	C1 for Yes with an example to illustrate	B		
	b	Yes The first equation has been divided by 2 throughout.	C1				C1 for correct language C1 for yes and a reason	
			3					
7	Substitute $x = 3$ in the equation to give $y = 3 + 2 = 5$ so when $x = 3$, $y = 5$, hence (2, 6) is not on the line or The constant term is 2 so the line crosses the y -axis at the point (0, 2). Then for every point across it goes up 1 (gradient is 1) so by the time $x = 3$, y will = 5.		C1	2	C1 for a clear example	B		
							1	
8	a	Is the sum of the cost of the CDs plus the coffee and the taxi less than £70?	C1	3	C1 for good question	B		
	b	Money spent = $2 \times £14.99 + 2 \times £2.50 + (12 \times £0.80 + £2.50)$ = $£29.98 + £5 + (£9.60 + £2.50)$ = $£34.98 + £12.10 = £47.08$ Money left = $£70 - £47.08 = £22.92$	Money left = $£70 - \text{money spent}$				B1	B1 for a correct formula that could be used
			This is less than £70 so she can afford the taxi.				P1	P1 for the process of calculating how much has been spent
							A1	A1 cao
							C1	C1 for clear, complete solution with correct answer
			5					
9	Look for the words that will represent variables and if possible, use appropriate letters to represent those variables. e.g. Area = height multiplied by breadth Formula could be $A = hb$		C1	2	C1 for an explanation of how to link a formula expressed in words to a formula expressed algebraically C1 if a suitable example has been included	B		
							2	
10	a	$2n$ means 2 times n while $n + 2$ means add 2 to n .	C1	2	C1 for clear explanation	B		
	b	$3(c + 5)$ means add 5 to c and then multiply the answer by 3, $3c + 5$ means multiply c by 3 and then add 5 to the answer.	C1				C1 for clear explanation	
	c	n^2 means multiply n by itself, $2n$ means multiply n by 2.	C1				C1 for clear explanation	
			3					

11	$\text{Perimeter} = 2 \times l + 2 \times 3l$ $= 2l + 6l = 8l$ So $8l = 48$ $l = 6 \text{ cm}$ $\text{Area} = \text{length} \times \text{width}$ $= l \times 3l$ $= 6 + 3 \times 6$ $= 6 + 18 = 24 \text{ cm}^2$	24 cm ²	M1 A1 M1 A1 4	3	M1 for using perimeter formula A1 cao M1 for area formula A1 ft	B
12	$\frac{(32 - 24)}{4} = 8 \div 4 = 2$ $24 - 2 \times 4 = 24 - 8$ $= 16$	C = 16	P1 A1 2	3	P1 for the correct process of working out C A1 cao	B
13		Plot the three points and draw the two sides. You can then complete the missing sides of the rectangle to complete the shape as shown in the diagram. Hence find the fourth vertex as in the diagram as (4, 8).	C1 C1 B1 3	2 3	C1 for clear explanation C1 for including a sketch alongside the explanation B1 for correctly indicating (4, 8)	B
14	Let the smaller number be n , then the next even number will be $(n + 2)$. $n + (n + 2) = 50$ $2n + 2 = 50$ $2n = 48$ $n = 24$ The lower number will be 24 so the larger number will be 26.	26	C1 M1 A1 A1 4	2 3	C1 for stating starting points M1 for method of setting up the equation A1 for solving for the first number A1 cao	B

15		<p>Example 1 As $24 = 6 \times 4$ $= 6 \times 2^2$ $t = ba^2$ Will give 24 when $b = 6$ and $a = 2$</p> <p>Example 2 As $24 = 3 \times 8$ $= 3 \times (2 + 6)$ $t = 3(a + b)$ Will give 24 when $a = 2$ and $b = 6$</p>	<p>B1 C1</p> <p>B1 C1</p> <p>4</p>	2	<p>B1 for first formula that works C1 for clear explanation of how it was found</p> <p>B1 for second formula that works C1 for clear explanation of how it was found</p>	M
16 a		<p>$5(c + 4) = 5c + 20$ Feedback 'Don't forget to multiply out both terms in the brackets.'</p>	<p>M1 C1</p>	2	<p>M1 for correctly expanding the brackets C1 for suitable feedback</p>	M
b		<p>$6(t - 2) = 6t - 12$ Feedback 'Don't forget 6(.....) means multiply both terms by 6.'</p>	<p>M1 C1</p>		<p>M1 for correctly expanding the brackets C1 for suitable feedback</p>	
c		<p>$-3(4 - s) = -12 + 3s$ Feedback 'Don't forget -3(.....) means multiply both terms by 6 and a minus x minus = ...</p>	<p>M1 C1</p>		<p>M1 for correctly expanding the brackets C1 for suitable feedback</p>	
d		<p>$15 - (n - 4) = 15 - n + 4 = 15 + 4 - n = 19 - n$ Feedback 'Don't forget - (n - 4) means multiply each term in the bracket by - 1 and that the - in the bracket belongs to the 4 to make it - 4.'</p>	<p>M1 C1</p> <p>8</p>		<p>M1 for correctly expanding the brackets C1 for suitable feedback</p>	
17		<p>Any equation in the form $y = mx + 1$ will pass through (0, 1) So $y = 2x + 1$ $y = 3x + 1$ will both pass through (0, 1)</p>	<p>C1</p> <p>B1 B1</p> <p>3</p>	2	<p>C1 for clear explanation</p> <p>B1 for first correct equation B1 for second correct equation</p>	M
18 a		<p>A correct example e.g. $2(z - 3) + 5q$</p>	<p>B1</p>	2	<p>B1 for an expression that is equivalent to $4z + 5q - 6$</p>	M
b		<p>A correct example e.g. $\frac{(10x - 4y)}{2}$</p>	<p>B1</p> <p>2</p>		<p>B1 for an expression that simplifies to $5x - 2y$</p>	

19 a		<table border="1"> <tbody> <tr> <td>x</td> <td>$12x$</td> <td>$2x$</td> </tr> <tr> <td>$7x$</td> <td></td> <td>$5x$</td> </tr> <tr> <td>$8x$</td> <td>$10x$</td> <td>$-2x$</td> </tr> </tbody> </table>	x	$12x$	$2x$	$7x$		$5x$	$8x$	$10x$	$-2x$	B4	3	B1 for each correct entry in the table	M
			x	$12x$	$2x$										
$7x$		$5x$													
$8x$	$10x$	$-2x$													
b	Own example that works.	P1 5	P1 for their own correct example that works												
20	$Z = 3A$ $Z = A + 18$ So $3A = A + 18$ $2A = 18$ $A = 9$ Substitute $A = 9$ into $Z = A + 18$ to give $Z = 27$ Check $3 \times 9 = 27$ which is correct.	Zoe has 27 and Alyssa has 9.	C1 C1 M1 A1 M1 A1	3	C1 for setting up first equation C1 for setting up second equation M1 for method of combining equations to eliminate one variable A1 for first correct answer found M1 for substituting first answer A1 for correct second answer	M									
21	$n + n + 20 = 2n + 20$ $2n + 20 = 90$ $2n = 70$ $n = 35$ So 35 on first shelf and $35 + 20 = 55$ on second. Need $90 \div 3 = 30$ on each shelf.	So need to move 5 from first shelf onto third shelf and 25 from second to third shelf.	C1 M1 A1 A1 P1	2 3	C1 for setting the initial expression M1 for setting this up to equal 90 A1 for first shelf as 35 A1 for second shelf as 55 P1 for correct process of sorting the books out to 30 on each shelf	M									
22		Select x values less than 0 and substitute into the equation.	P1	2	P1 for clear explanation	M									
			1												

27	<p>Sketch a graph:</p> 	11 am	C1 C1 C1 B1 4	3	C1 for showing runner on graph or explaining C1 for showing cyclist on graph or explaining C1 for showing where the two lines meet on graph or explaining B1 cao	M
28	100, 96, 92, 88, 84, 80, 76, 72, 68, 64, 60, 56, 52, 48, 44, 40, 36, 32, 28, 24, 20, 16, 12, 8, 4 2, 8, 14, 20, 26, 32, 38, 44, 50, 56, 62, 68, 74, 80, 86, 92, 98 Those in common 8, 20, 32, 44, 56, 68, 80, 92	8, 20, 32, 44, 56, 68, 80, 92	P1 P1 B1 3	3	P1 for process of accounting for first sequence P1 for process of accounting for second sequence B1 for all 8 correct terms	M
29	Left hand graph is $x + y = 5$ Right hand graph is $y = z + 1$ Substitute y into first equation $x + z + 1 = 5$ $x + z = 4$		B1 B1 M1 A1 C2 6	3	B1 first graph equation B1 second graph equation M1 substituting to eliminate y A1 cao C1 for graph drawn with x on vertical axis. Allow x on horizontal axis C1 for $x + z = 4$ drawn correctly	M
30 a	Distance = $2 \times 25 \text{ km} = 50 \text{ km}$ 50 km \div 8 hours = 6.25 km per hour.	6.25 km/h	M1 A1 C1 C2 5	2 3	M1 for division of total distance by time A1 cao C1 for an example of a questions that could be asked about this situation C1 for a two part question using the graph with increase in difficulty C1 for suitable mark scheme	M
31		Own story Sketch graph Question for the graph	C1 C1 C1 3	2	C1 for suitable story C1 for matching sketch graph C1 for suitable question	M

32	a i	$35 \times 8 + 10$	£290	M1 A1	3	M1 for the correct method A1 cao	M
	ii	35×14	£490	M1 A1		M1 for correct method A1 cao	
	b	$35n + 10 = 220$ $35n = 210$ $n = \frac{210}{35} = 6$	6 sessions	P1 A1		P1 for process of sorting which rule to use A1 cao	
	c	$(7 \times 35) + 20 = \text{£}265$ $(7 \times 35) = 10 = \text{£}255$	£10 more	P1 A1		P1 for finding suitable calculations to find the difference A1 cao	
				8			
33		$10 + 15 = 25 = 5^2$ $15 + 21 = 36 = 6^2$		B4 4		B1 for each correct part of the number pattern provided correct signs and symbols are present	M
34	a		Triangle drawn	C1	3	C1 for diagram drawn for all shapes	M
	b		36 cm	B1		B1 cao	
	c		48 cm	B1		B1 cao	
	d / e		63 139 143 806 710 cm	B1 4		B1 cao	
35	a		Same difference of 2.4 but starting value is different.	B1 B1	2	B1 cao B1 cao	M
	b		What are the differences What is the starting value.	B1 B1 4		B1 cao B1 cao	
36	a		Multiple of 4	B1 B1 C1	2	B1 cao	M
	b		No because we need to know the starting value as well.	3		C1 for no C1 for reason alongside no	

37		<p>Boys Get a red egg each from each of 4 girls: 4 red One green egg each other: 2 green</p> <p>Girls Get a blue egg from each of the 2 boys: 2 blue One yellow egg from each other will be 3 yellow eggs each: 12 yellow</p>	<p>C1 C1 C1 C1 5</p>	3	<p>C1 for explanation of 4 red C1 for explanation of 2 green C1 for explanation of 2 blue C1 for explanation of 12 yellow C1 for complete clear solution</p>	M
38	<p>Example $2n^2 = 2 \times (3^2) = 2 \times 9 = 18$ $(2 \times 3)^2 = 6 \times 6 = 36$</p>	<p>Using BIDMAS for $2n^2$ tells you to calculate the power first. BIDMAS for $(2n)^2$ tells you that you do the calculation inside the bracket first.</p>	<p>C1 5</p>	2	<p>C1 for an explanation. An example could be given to support the argument</p>	M
39		<p>A letter, say f, stands for an unknown if it is in an equation such as $3f + 2 = 14$. Then $f = 4$ is the only number that satisfies this equation.</p> <p>A letter stands for a variable if it is part of an equation that has more than two letters. E.g. $A = \pi r^2$, where both A and r are variables that will be different for different values of A or r.</p>	<p>C1 C1 C1 C1 4</p>	2	<p>C1 for clear explanation C1 for an example alongside the explanation C1 for a clear explanation C1 for an example alongside the explanation</p>	M

42		$\frac{(2n+6)}{2}$ $= \frac{2(n+3)}{2} = n+3$	M1 B1 2	2	M1 for factorising B1 for any correct expression	M
43		<p>Let base length be b, then height will be $3b$</p> <p>Area of triangle = $\frac{1}{2} \times \text{base} \times \text{height}$</p> $= \frac{1}{2} \times b \times 3b$ $= \frac{3}{2}b^2$ <p>Where $A = 6$</p> $\frac{1}{2}b^2 = 6$ $b^2 = 2 \times \frac{6}{3} = 4$ $b = 2$ <p>so height is 3×2 which is 6 cm.</p>	C1 C1 B1 M1 A1 A1 6	3	C1 for stating variables C1 for stating triangle formula B1 for correct expression M1 for equating 6 with found expression A1 for $b = 2$ A1 for 6 cm	M

<p>44 a</p> <p>b</p>	<table border="1" data-bbox="264 185 656 347"> <thead> <tr> <th>x</th> <th>x^3</th> <th>$x + x^3$</th> <th>Too...</th> </tr> </thead> <tbody> <tr><td>1</td><td>1</td><td>2</td><td>small</td></tr> <tr><td>2</td><td>8</td><td>10</td><td>small</td></tr> <tr><td>3</td><td>27</td><td>30</td><td>big</td></tr> <tr><td>2.5</td><td>15.63</td><td>18.13</td><td>small</td></tr> <tr><td>2.6</td><td>17.58</td><td>20.18</td><td>big</td></tr> <tr><td>2.55</td><td>16.58</td><td>19.13</td><td>small</td></tr> </tbody> </table> <table border="1" data-bbox="264 373 656 491"> <thead> <tr> <th>x</th> <th>$x + 2$</th> <th>$x(x + 2)$</th> <th>Too...</th> </tr> </thead> <tbody> <tr><td>7</td><td>9</td><td>63</td><td>small</td></tr> <tr><td>8</td><td>10</td><td>80</td><td>big</td></tr> <tr><td>7.5</td><td>9.5</td><td>71.25</td><td>big</td></tr> <tr><td>7.3</td><td>9.3</td><td>67.89</td><td>exact</td></tr> </tbody> </table>	x	x^3	$x + x^3$	Too...	1	1	2	small	2	8	10	small	3	27	30	big	2.5	15.63	18.13	small	2.6	17.58	20.18	big	2.55	16.58	19.13	small	x	$x + 2$	$x(x + 2)$	Too...	7	9	63	small	8	10	80	big	7.5	9.5	71.25	big	7.3	9.3	67.89	exact	<p>You could use trial and improvement or a graph to help you decide where to start.</p> <p>Use trial and improvement to solve both problems.</p> <p>Number is 2.6</p> <p>Width is 7.3 cm</p>	<p>C1</p> <p>P1 P2</p> <p>A1</p> <p>P2</p> <p>A1</p> <p>8</p>	<p>2</p>	<p>C1 for explanation of suitable methods, could also be graphs</p> <p>P1 for using their suggested method(s) P1 for finding the range including the solution P1 for process of finding which of the 1 dp trials is closest</p> <p>A1 for 2.6 or more accurate</p> <p>P1 for finding the range including the solution P1 for process of finding which of the 1 dp trials is closest</p> <p>A1 cao</p>	<p>H</p>
x	x^3	$x + x^3$	Too...																																																			
1	1	2	small																																																			
2	8	10	small																																																			
3	27	30	big																																																			
2.5	15.63	18.13	small																																																			
2.6	17.58	20.18	big																																																			
2.55	16.58	19.13	small																																																			
x	$x + 2$	$x(x + 2)$	Too...																																																			
7	9	63	small																																																			
8	10	80	big																																																			
7.5	9.5	71.25	big																																																			
7.3	9.3	67.89	exact																																																			
<p>45 a</p> <p>b i</p> <p>ii</p> <p>iii</p>		<p>'I think of a number and double it' just has an expression of $2x$ where x is the number I thought of – still unknown at the moment.</p> <p>'I think of a number and double it – the answer is 12' has a solution that I know is 6.</p> <p>One</p> <p>e.g. $10 = p + 3$</p> <p>Because each solution is $p = 7$</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>4</p>	<p>2</p>	<p>B1 for clear explanation of the difference</p> <p>B1 cao</p> <p>B1 for a correct example</p> <p>B1 for a clear explanation</p>	<p>H</p>																																																

46	<p>Looking at total counters needed for each step, he uses: Step 1: 6 counters Step 2: 12 counters Step 3: 18 counters Step 4: 24 counters Step n: $6n$ counters</p> <p>Adding how many counters he needs in total: Step 1: 6 counters Step 2: 18 counters Step 3: 36 counters Step 4: 60 counters</p> <p>Looking at the pattern suggests products being involved, I see that this pattern can be written as Step 1: $3 \times 1 \times 2 = 6$ Step 2: $3 \times 2 \times 3 = 18$ Step 3: $3 \times 3 \times 4 = 36$ Step 4: $3 \times 4 \times 5 = 60$ Step n: $3n(n + 1)$</p> <p>I need to find a value for n where this total</p> <table border="1"> <thead> <tr> <th>n</th> <th>$n + 1$</th> <th>$3n(n + 1)$</th> <th>Too...</th> </tr> </thead> <tbody> <tr> <td>10</td> <td>11</td> <td>330</td> <td>small</td> </tr> <tr> <td>20</td> <td>21</td> <td>1260</td> <td>big</td> </tr> <tr> <td>15</td> <td>16</td> <td>720</td> <td>small</td> </tr> <tr> <td>17</td> <td>18</td> <td>918</td> <td>small</td> </tr> <tr> <td>18</td> <td>19</td> <td>1026</td> <td>big</td> </tr> </tbody> </table> <p>is first over 1000 Use trial and improvement</p>	n	$n + 1$	$3n(n + 1)$	Too...	10	11	330	small	20	21	1260	big	15	16	720	small	17	18	918	small	18	19	1026	big	<p>Harry will run out of counters while trying to complete step 18.</p>	B1	3	<p>B1 for the process of finding how many counters needed for each step</p> <p>P1 for $6n$</p> <p>P1 for the process of finding the total number of counters used by each step</p> <p>B1 for the process of looking to generalise this pattern</p> <p>C1</p> <p>B1 for the generalisation</p> <p>P1 for the explanation of what he needed to do. A1 for a suitable process of finding which step he would get to</p> <p>A1</p> <p>A1 cao</p>	H
		n	$n + 1$	$3n(n + 1)$	Too...																									
10	11	330	small																											
20	21	1260	big																											
15	16	720	small																											
17	18	918	small																											
18	19	1026	big																											
P1	8																													
47		No. All the terms will be even.	B1	2	<p>B1 for no C1 for clear explanation</p>	H																								
			C1				2																							

50	a	1 6 15 20 15 6 1 1 7 21 35 35 21 7 1 1 8 28 56 70 56 28 8 1		B1	2 3	B1 for correct next three rows	H																											
	b	Looking at the diagonal rows: The first diagonal row contains only 1s. The second diagonal consists of all counting numbers: 1, 2, 3, 4, 5, etc. The third row consists of the triangle numbers: 1, 3, 6, 10, 15, etc.		C3																														
	c	Triangle numbers		B1	B1 for triangle numbers																													
	d	1, 2, 4, 8, 16, 32		B1	B1 for correct sequence																													
	e	Multiplying by 2 each time, the n th term will be 2^{n-1}		C1	C1 for clear explanation																													
				7																														
51		$6(x - c) = 5x - 4$ $6x - 6c = 5x - 4$ $x = 6c - 4$ $6c$ is always even as even \times odd/even = even 4 is even So x must be even as even $-$ even = even		M1	2	M1 for expanding the bracket	H																											
				A1				A1 for x as subject																										
				C1				C1 for clear explanation																										
				3																														
52	<table border="1"> <thead> <tr> <th>n</th> <th>1</th> <th>2</th> <th>3</th> <th>4</th> <th>5</th> <th>6</th> <th>7</th> <th>8</th> </tr> </thead> <tbody> <tr> <td>$\frac{n+1}{2n+1}$</td> <td>$\frac{2}{3}$</td> <td>$\frac{3}{5}$</td> <td>$\frac{4}{7}$</td> <td>$\frac{5}{9}$</td> <td>$\frac{6}{11}$</td> <td>$\frac{7}{13}$</td> <td>$\frac{8}{15}$</td> <td>$\frac{9}{17}$</td> </tr> <tr> <td></td> <td>0.66̄</td> <td>0.6</td> <td>0.571428̄</td> <td>0.55</td> <td>0.54</td> <td>0.538461̄</td> <td>0.533</td> <td>0.52941176̄</td> </tr> </tbody> </table>		n	1	2	3	4	5	6	7	8	$\frac{n+1}{2n+1}$	$\frac{2}{3}$	$\frac{3}{5}$	$\frac{4}{7}$	$\frac{5}{9}$	$\frac{6}{11}$	$\frac{7}{13}$	$\frac{8}{15}$	$\frac{9}{17}$		0.66̄	0.6	0.571428̄	0.55	0.54	0.538461̄	0.533	0.52941176̄		C1	2	C1 for showing the pattern of fractions	H
	n	1	2	3	4	5	6	7	8																									
	$\frac{n+1}{2n+1}$	$\frac{2}{3}$	$\frac{3}{5}$	$\frac{4}{7}$	$\frac{5}{9}$	$\frac{6}{11}$	$\frac{7}{13}$	$\frac{8}{15}$	$\frac{9}{17}$																									
		0.66̄	0.6	0.571428̄	0.55	0.54	0.538461̄	0.533	0.52941176̄																									
			C1	C1 for showing all the decimals																														
Only $\frac{3}{5}$ is a terminating decimal.			C1	C1 for clear explanation																														
				3																														
53	a		even	B1	2	B1 cao	H																											
	b		odd	B1																														
	c		even	B1																														
	d		even	B1																														
	e		even	B1																														
	f		even	B1																														
	g		odd	B1																														
	h		even	B1																														
				8																														

54	a	$t = \frac{6}{n}$: Graph B One person will take a long time, many people will take a short time.	B2 C1	2 3	B1 for correct equation B1 for correct graph C1 for good reason for choice	H
	b	$s = -4.9t^2 + 40t + 80$: Graph D This is a quadratic graph and it shows the value 80 when t is 0, the height of the cliff.	B2 C1			
	c	$y = 3x + 320$: Graph A This will be a linear graph and this graph also crosses the vertical axis at (320, 0) showing his starting pay before selling any items.	B2 C1			
	d	$x^2 + 72x - 225 = 0$: Graph C The area from the dimensions will create a quadratic graph which moves further and further into the first quadrant.	B2 C1			
			12			
55		<p>c and d can be difficult because they contain minus signs and this is a point where errors are made, combining minus signs.</p> <p>In substituting $x = -3$ into $t = -2(3 - x)$, a classic error is to assume $3 - -3$ is 0.</p> <p>In substituting $x = -3$ into $z = \frac{-2(x + 2)}{x}$, a classic error is to give a negative divided by a negative a negative answer.</p> <p>A suggestion to avoid these errors is to remember that when multiplying or dividing with positive and negative numbers, same signs means positive, different signs means negative.</p>	C1 C2 C1	2	C1 for identifying some examples with a valid reason C1 for clear identification of one classic error with one equation C1 for another classic error C1 for a satisfactory suggestion	H
			4			

56		The similarities are that both have an equals sign and both require the manipulation of terms.	C1	2	C1 for clear explanation of similarities	H
		The difference is that in solving an equation you end up with a numerical answer, but in rearranging you still have a formula.	C1		C1 for clear explanation of differences	
57	a	The two straight-line graphs will be parallel, with the same gradient of 2. $y = 2x$ crosses the y -axis at the origin and $y = 2x + 6$ crosses the y -axis at $y = 6$	C2	2	C1 for explanation of parallel C1 for explanation containing points of intersection of axes	H
	b	The two straight-line graphs will be parallel, with the same gradient of 1. $y = x + 5$ crosses the y -axis at $y = 5$, and $y = x - 6$ crosses the y -axis at $y = -6$	C2		C1 for explanation of parallel C1 for explanation containing points of intersection of axes	
	c	The two straight-line graphs will cross each other at $(\frac{11}{8}, \frac{1}{2})$ and each one is a reflection of the other in a vertical mirror line.	C2		C1 for explanation containing point of intersection C1 for explanation of symmetry	
	d	The two straight-line graphs will both cross the y -axis at the origin, one with gradient 2, another with a gradient of $\frac{1}{2}$.	C2		C1 for explanation of passing through origin C1 for explanation about gradient	
			8			